

Anti-MCH Receptor 1 Antibody
Catalog # AP54085**Specification**

Anti-MCH Receptor 1 Antibody - Product Information

Application	WB, IF
Primary Accession	Q969V1
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	38849

Anti-MCH Receptor 1 Antibody - Additional Information**Gene ID** 84539**Other Names**GPR145; SLT; Melanin-concentrating hormone receptor 2; MCH receptor 2; MCH-R2; MCHR-2;
G-protein coupled receptor 145; GPRv17; MCH-2R; MCH2; MCH2R**Target/Specificity**

Recognizes endogenous levels of MCH Receptor 1 protein.

Dilution

WB~~1/500 - 1/1000

IF~~1/50 - 1/200

Format

Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.

Storage

Store at -20 °C.Stable for 12 months from date of receipt

Anti-MCH Receptor 1 Antibody - Protein Information**Name** MCHR2**Synonyms** GPR145, SLT**Function**

Receptor for melanin-concentrating hormone, coupled to G proteins that activate phosphoinositide hydrolysis.

Cellular Location

Cell membrane; Multi-pass membrane protein.

Tissue Location


Specifically expressed in the brain, with highest levels in cerebral cortex, hippocampus and amygdala. No expression detected in the cerebellum, thalamus or hypothalamus

Anti-MCH Receptor 1 Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-MCH Receptor 1 Antibody - Images



Western blot analysis of MCH Receptor 1 expression in rat testis (A), BV2 (B), LOVO (C), HCT116 (D) whole cell lysates.



Immunofluorescent analysis of MCH Receptor 1 staining in LOVO cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a Alexa Fluor 594-conjugated secondary antibody (red) in PBS at room temperature in the

dark.

Anti-MCH Receptor 1 Antibody - Background

Rabbit polyclonal antibody to MCH Receptor 1