

NPSR1 Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP54498**Specification****NPSR1 Polyclonal Antibody - Product Information**

Application	WB, IHC-P, IHC-F, IF, ICC, E
Primary Accession	Q6W5P4
Reactivity	Rat, Pig, Dog, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	42687

NPSR1 Polyclonal Antibody - Additional Information**Gene ID** 387129**Other Names**

Neuropeptide S receptor, G-protein coupled receptor 154, G-protein coupled receptor PGR14, G-protein coupled receptor for asthma susceptibility, NPSR1, GPR154, GPRA, PGR14

Dilution

WB~~1:1000<br \>IHC-P~~N/A<br \>IHC-F~~N/A<br \>IF~~1:50~200<br \>ICC~~N/A<br \>E~~N/A

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

NPSR1 Polyclonal Antibody - Protein Information**Name** NPSR1**Synonyms** GPR154, GPRA, PGR14**Function**

G-protein coupled receptor for neuropeptide S (NPS) (PubMed:16790440). Promotes mobilization of intracellular Ca(2+) stores (PubMed:16790440). Inhibits cell growth in response to NPS binding (PubMed:15947423). Involved in pathogenesis of asthma and other IgE- mediated

diseases.

Cellular Location

[Isoform 1]: Cell membrane; Multi-pass membrane protein [Isoform 4]: Cell membrane; Multi-pass membrane protein [Isoform 5]: Cytoplasm [Isoform 7]: Cytoplasm

Tissue Location

Isoform 4 is ubiquitous; it is detected in glandular epithelia of bronchus, stomach, small intestine, colon, uterus, esophagus, spleen, kidney, pancreas, prostate and breast Isoform 1 is detected in uterus, colon and prostate, and in the smooth muscle cell layer in bronchial and arterial walls (at protein level) (PubMed:15947423). Isoform 1 is predominantly expressed in smooth muscle.

Isoform 4 is predominantly expressed in epithelial cells. In bronchial biopsies, it is expressed in smooth muscle cells of asthma patients, but not in control patients; whereas in epithelial cells, its expression is consistently stronger in asthma patients

NPSR1 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

NPSR1 Polyclonal Antibody - Images