

#### **SMPDL3B Polyclonal Antibody**

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP54917

#### **Specification**

# **SMPDL3B Polyclonal Antibody - Product Information**

Application IHC-P, IHC-F, IF, ICC, E

Primary Accession
Reactivity
Rat
Host
Clonality
Calculated MW
Physical State

Q92485
Rat
Rabbit
Polyclonal
49 KDa
Liquid

Immunogen KLH conjugated synthetic peptide derived

from human SMPDL3B

Epitope Specificity 181-280/455

lsotype IgG

Purity
affinity purified by Protein A

Buffer 0.01M TBS (pH7.4) with 1% BSA, 0.02%

Proclin300 and 50% Glycerol.

SUBCELLULAR LOCATION Secreted.

SIMILARITY Belongs to the acid sphingomyelinase

family

SUBUNIT Interacts with TLR4, TLR7, TLR8 and TLR9.

Important Note This product as supplied is intended for

research use only, not for use in human, therapeutic or diagnostic applications.

# **Background Descriptions**

Located on chromosome 1, this gene encodes for acid sphingomyelinase like phosphodiesterase 3b precursor protein.

# **SMPDL3B Polyclonal Antibody - Additional Information**

# **Gene ID 27293**

# **Other Names**

Acid sphingomyelinase-like phosphodiesterase 3b, ASM-like phosphodiesterase 3b, 3.1.4.-, SMPDL3B, ASML3B, ASMLPD

# **Dilution**

<span class ="dilution IHC-P">IHC-P~~N/A</span><br \><span class</pre>

="dilution IHC-F">IHC-F~~N/A</span><br \><span class

="dilution\_IF">IF~~1:50~200</span><br \><span class = "dilution\_ICC">ICC~~N/A</span><br \><span class = "dilution\_E">E~~N/A</span>

#### **Format**

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce



**Storage** 

Store at -20  $^{\circ}$ C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4  $^{\circ}$ C.

# **SMPDL3B Polyclonal Antibody - Protein Information**

Name SMPDL3B

Synonyms ASML3B, ASMLPD

#### **Function**

Lipid-modulating phosphodiesterase (PubMed:<a

href="http://www.uniprot.org/citations/26095358" target="\_blank">26095358</a>). Active on the surface of macrophages and dendritic cells and strongly influences macrophage lipid composition and membrane fluidity. Acts as a negative regulator of Toll-like receptor signaling (By similarity). Has in vitro phosphodiesterase activity, but the physiological substrate is unknown (PubMed:<a href="http://www.uniprot.org/citations/26095358" target="\_blank">26095358</a>). Lacks activity with phosphocholine-containing lipids, but can cleave CDP-choline, and can release phosphate from ATP and ADP (in vitro) (By similarity).

#### **Cellular Location**

Secreted. Cell membrane  $\{ECO:0000250|UniProtKB:P58242\}$ ; Lipid-anchor, GPI-anchor  $\{ECO:0000250|UniProtKB:P58242\}$ 

# **SMPDL3B Polyclonal Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

#### **SMPDL3B Polyclonal Antibody - Images**