

SYBL1 Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP55002

Specification

SYBL1 Polyclonal Antibody - Product Information

Application WB, IHC-P, IHC-F, IF, ICC

Primary Accession <u>P51809</u>

Reactivity Rat, Pig, Bovine

Host Rabbit
Clonality Polyclonal
Calculated MW 25 KDa
Physical State Liquid

Immunogen KLH conjugated synthetic peptide derived

from human SYBL1

Epitope Specificity 1-100/220

Isotype IgG
Purity

affinity purified by Protein A

Buffer 0.01M TBS (pH7.4) with 1% BSA, 0.02%

Proclin300 and 50% Glycerol.

SUBCELLULAR LOCATION Cytoplasmic vesicle > secretory vesicle

membrane. Golgi apparatus > trans-Golgi network membrane. Late endosome membrane. Lysosome membrane. Endoplasmic reticulum membrane. Cytoplasmic vesicle > phagosome

membrane.

SIMILARITY Belongs to the synaptobrevin family.

Contains 1 longin domain. Contains 1 v-SNARE coiled-coil homology domain. This product as supplied is intended for research use only, not for use in human,

therapeutic or diagnostic applications.

Background Descriptions

FLJ53045; FLJ53762; FLJ54296; HGNC:11486; OTTHUMP00000024258; OTTHUMP00000024259; OTTHUMP00000225953; SYBL 1; SYBL1; Synaptobrevin like 1; Synaptobrevin-like protein 1; Tetanus insensitive VAMP; Tetanus neurotoxin insensitive VAMP; Tetanus-insensitive VAMP; TIVAMP; TIVAMP; VAMP-7; VAMP7; VAMP7 HUMAN

SYBL1 Polyclonal Antibody - Additional Information

Gene ID 6845

Important Note

Other Names

Vesicle-associated membrane protein 7, VAMP-7, Synaptobrevin-like protein 1, Tetanus-insensitive VAMP, Ti-VAMP, VAMP7, SYBL1



Target/Specificity

Detected in all tissues tested.

Dilution

- WB~~1:1000<br \><span class</pre>
- ="dilution IHC-P">IHC-P~~N/A<br \><span class
- ="dilution IHC-F">IHC-F~~N/A<br \><span class
- ="dilution_IF">IF \sim 1:50 \sim 200<br\>ICC \sim N/A

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

SYBL1 Polyclonal Antibody - Protein Information

Name VAMP7

Synonyms SYBL1

Function

Involved in the targeting and/or fusion of transport vesicles to their target membrane during transport of proteins from the early endosome to the lysosome. Required for heterotypic fusion of late endosomes with lysosomes and homotypic lysosomal fusion. Required for calcium regulated lysosomal exocytosis. Involved in the export of chylomicrons from the endoplasmic reticulum to the cis Golgi. Required for exocytosis of mediators during eosinophil and neutrophil degranulation, and target cell killing by natural killer cells. Required for focal exocytosis of late endocytic vesicles during phagosome formation.

Cellular Location

Cytoplasmic vesicle, secretory vesicle membrane; Single-pass type IV membrane protein Golgi apparatus, trans-Golgi network membrane; Single- pass type IV membrane protein. Late endosome membrane; Single-pass type IV membrane protein Lysosome membrane; Single-pass type IV membrane protein. Endoplasmic reticulum membrane; Single-pass type IV membrane protein. Cytoplasmic vesicle, phagosome membrane; Single-pass type IV membrane protein. Synapse, synaptosome. Note=In immature neurons expression is localized in vesicular structures in axons and dendrites while in mature neurons it is localized to the somatodendritic region Colocalizes with LAMP1 in kidney cells. Localization to the endoplasmic reticulum membrane was observed in the intestine but not in liver or kidney (By similarity).

Tissue Location

Detected in all tissues tested.

SYBL1 Polyclonal Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry



- <u>Immunofluorescence</u>
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

SYBL1 Polyclonal Antibody - Images



Sample:

Liver (Mouse) Lysate at 40 ug

Primary: Anti-SYBL1 (bs-12852R) at 1/300 dilution

Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution

Predicted band size: 25 kD Observed band size: 26 kD