

Vinexin Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP55265

Specification

Vinexin Polyclonal Antibody - Product Information

Application WB, IHC-P, IHC-F, IF, ICC, E

Primary Accession <u>060504</u>

Reactivity Rat, Pig, Dog, Bovine

Host Rabbit
Clonality Polyclonal
Calculated MW 75341

Vinexin Polyclonal Antibody - Additional Information

Gene ID 10174

Other Names

Vinexin, SH3-containing adapter molecule 1, SCAM-1, Sorbin and SH3 domain-containing protein 3, SORBS3, SCAM1

Dilution

WB~~1:1000<br \><span class
="dilution_IHC-P">IHC-P~~N/A<br \><span class
="dilution_IHC-F">IHC-F~~N/A<br \><span class
="dilution_IF">IF~~1:50~200<br \>ICC~~N/A<br \>ICC~~N/A<br \>ICC~~N/A

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Vinexin Polyclonal Antibody - Protein Information

Name SORBS3

Synonyms SCAM1

Function

Vinexin alpha isoform promotes up-regulation of actin stress fiber formation. Vinexin beta isoform plays a role in cell spreading and enhances the activation of JNK/SAPK in response to EGF stimulation by using its third SH3 domain.

Cellular Location

[Isoform Alpha]: Cell junction. Cytoplasm, cytoskeleton. Note=Localized at cell- extracellular



matrix junctions (By similarity). Both isoforms were localized at focal adhesion and cell-cell adhesion sites

Tissue Location

Both isoforms are expressed in different tissues like heart, placenta, brain, skeletal muscle and pancreas. Isoform beta is especially found in liver

Vinexin Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

Vinexin Polyclonal Antibody - Images