

Pannexin 1 Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP57823

Specification

Pannexin 1 Polyclonal Antibody - Product Information

Application WB, IHC-P, IHC-F, IF, ICC, E

Primary Accession
Reactivity
Rat, Bovine
Host
Clonality
Calculated MW
Rat, Bovine
Rabbit
Polyclonal
48 KDa

Physical State
Liquid
Immunogen
KLH conjugated synthetic peptide derived

from human Pannexin 1

Epitope Specificity 61-106/426

Isotype IgG

affinity purified by Protein A

Buffer 0.01M TBS (pH7.4) with 1% BSA, 0.02%

Proclin300 and 50% Glycerol.

SUBCELLULAR LOCATION Cell membrane. Cell junction > gap

junction. Endoplasmic reticulum

membrane.

SIMILARITY
SUBUNIT

Belongs to the pannexin family.
Homohexamer. Forms homomeric or

PANX1/PANX2-heteromeric intercellular channels on coexpression in paired Xenopus oocytes (By similarity).

Important Note

This product as supplied is intended for

research use only, not for use in human, therapeutic or diagnostic applications.

Background Descriptions

The protein encoded by this gene belongs to the innexin family. Innexin family members are the structural components of gap junctions. This protein and pannexin 2 are abundantly expressed in central nerve system (CNS) and are coexpressed in various neuronal populations. Studies in Xenopus oocytes suggest that this protein alone and in combination with pannexin 2 may form cell type-specific gap junctions with distinct properties. [provided by RefSeq, Jul 2008]

Pannexin 1 Polyclonal Antibody - Additional Information

Gene ID 24145

Other Names

Pannexin-1. PANX1 (<a

href="http://www.genenames.org/cgi-bin/gene symbol report?hgnc id=8599"

target="_blank">HGNC:8599), MRS1



Dilution

```
 < span \ class = "dilution_WB">WB~\sim 1:1000 < /span> < br \> < span \ class = "dilution_IHC-P">IHC-P~\sim N/A < /span> < br \> < span \ class = "dilution_IHC-F">IHC-F~\sim N/A < /span> < br \> < span \ class = "dilution_IF">IF~\sim 1:50 \sim 200 < /span> < br \> < span \ class = "dilution_ICC">ICC~\sim N/A < /span> < br \> < span \ class = "dilution_E">E~\sim N/A < /span> < br \> < span \ class = "dilution_E">E~\sim N/A < /span> < br \> < span \ class = "dilution_E">E~\sim N/A < /span> < br \> < span \ class = "dilution_E">E~\sim N/A < /span> < br \> < span \ class = "dilution_E">E~\sim N/A < /span> < br \> < span \ class = "dilution_E">E~\sim N/A < /span> < br \> < span \ class = "dilution_E">E~\sim N/A < /span> < br \> < span \ class = "dilution_E">E~\sim N/A < /span> < br \> < span \ class = "dilution_E">E~\sim N/A < /span> < br \> < span \ class = "dilution_E">E~\sim N/A < /span> < br \> < span \ class = "dilution_E">E~\sim N/A < /span> < br \> < span \ class = "dilution_E">E~< \ D/A < /span < down = 10000 < down = 100000 < down = 10000 < down = 100000 < down = 10000 < down = 100000 < down =
```

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 $^{\circ}$ C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 $^{\circ}$ C.

Pannexin 1 Polyclonal Antibody - Protein Information

Name PANX1 (HGNC:8599)

Function

Ion channel involved in a variety of physiological functions such as blood pressure regulation, apoptotic cell clearance and oogenesis (PubMed:15304325, PubMed:16908669, PubMed:20829356, PubMed:20944749, PubMed:30918116). Forms anion-selective channels with relatively low conductance and an order of permeabilities: nitrate>iodide>chlroride>>aspartate=glutamate=gluconate (By similarity). Can release ATP upon activation through phosphorylation or cleavage at C-terminus (PubMed:32238926). May play a release a Ca(2+) look channel to regulate ER Ca(2+) homeostasis (PubMed:<a

role as a Ca(2+)- leak channel to regulate ER Ca(2+) homeostasis (PubMed:16908669).

Cellular Location

Cell membrane; Multi-pass membrane protein {ECO:0000255|PROSITE-ProRule:PRU00351}. Endoplasmic reticulum membrane; Multi-pass membrane protein {ECO:0000255|PROSITE-ProRule:PRU00351}

Tissue Location

Widely expressed (PubMed:30918116). Highest expression is observed in oocytes and brain (PubMed:30918116). Detected at very low levels in sperm cells (PubMed:30918116)

Pannexin 1 Polyclonal Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety



• Cell Culture

Pannexin 1 Polyclonal Antibody - Images



Sample: Muscle (Mouse) Lysate at 40 ug

Primary: Anti-Pannexin 1(bs-21015R)at 1/300 dilution

Secondary: IRDye800CW Goat Anti-RabbitlgG at 1/20000 dilution

Predicted band size: 48kD Observed band size: 53kD