

KDEL Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP58558**Specification****KDEL Polyclonal Antibody - Product Information**

Application	IHC-P, IHC-F, IF, E
Primary Accession	Q6UW63
Reactivity	Rat, Pig, Dog, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	78 KDa
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide KDEL
Isotype	IgG
Purity	
affinity purified by Protein A	
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Endoplasmic reticulum lumen.
SIMILARITY	Belongs to the KDELC family. Contains 1 filamin repeat.
SUBUNIT	Interacts with DNAJC1 (via J domain). Component of an EIF2 complex at least composed of CELF1/CUGBP1, CALR, CALR3, EIF2S1, EIF2S2, HSP90B1 and HSPA5. Part a large chaperone multiprotein complex comprising DNAJB11, HSP90B1, HSPA5, HYOU, PDIA2, PDIA4, PDIA6, PPIB, SDF2L1, UGT1A1 and very small amounts of ERP29, but not, or at very low levels, CALR nor CANX. Interacts with TMEM132A and TRIM21. May form a complex with ERLEC1, OS9, SEL1L and SYVN1.
Post-translational modifications	N-glycosylated.
DISEASE	Note=Autoantigen in rheumatoid arthritis.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Background Descriptions

This gene encodes a protein product localized to the lumen of the endoplasmic reticulum. As a member of the endoplasmic reticulum protein family the encoded protein contains a Lys-Asp-Glu-Leu or KDEL motif located at the extreme C-terminus which prevents all endoplasmic reticulum resident proteins from being secreted. Proteins carrying this motif are bound by a receptor in the Golgi apparatus so that the receptor-ligand complex returns to the endoplasmic reticulum. A processed non-transcribed pseudogene located in an intron of a sodium transporter gene on chromosome 5 has been defined for this gene. [provided by RefSeq, Jul 2008]

KDEL Polyclonal Antibody - Additional Information

Gene ID 79070**Other Names**

Protein O-glucosyltransferase 2, 2.4.1.-, Endoplasmic reticulum resident protein 58, ER protein 58, ERp58, KDEL motif-containing protein 1 {ECO:0000312|HGNC:HGNC:19350}, Protein O-xylosyltransferase POGLUT2, 2.4.2.-, POGLUT2 {ECO:0000303|PubMed:30127001, ECO:0000312|HGNC:HGNC:19350}

Dilution

IHC-P ~ N/A
IHC-F ~ N/A
IF ~ 1:50 ~ 200
E ~ N/A

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

KDEL Polyclonal Antibody - Protein Information

Name POGLUT2 {ECO:0000303|PubMed:30127001, ECO:0000312|HGNC:HGNC:19350}

Function

Protein glucosyltransferase that catalyzes the transfer of glucose from UDP-glucose to a serine residue within the consensus sequence peptide C-X-N-T-X-G-S-F-X-C (PubMed: [30127001](http://www.uniprot.org/citations/30127001)). Can also catalyze the transfer of xylose from UDP-xylose but less efficiently (PubMed: [30127001](http://www.uniprot.org/citations/30127001)). Specifically targets extracellular EGF repeats of proteins such as NOTCH1, NOTCH3, FBN1, FBN2 and LTBP1 (PubMed: [30127001](http://www.uniprot.org/citations/30127001), PubMed: [34411563](http://www.uniprot.org/citations/34411563)). May regulate the transport of NOTCH1 and NOTCH3 to the plasma membrane and thereby the Notch signaling pathway (PubMed: [30127001](http://www.uniprot.org/citations/30127001)).

Cellular Location


Endoplasmic reticulum lumen {ECO:0000255|PROSITE- ProRule:PRU10138}

KDEL Polyclonal Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

KDEL Polyclonal Antibody - Images



Paraformaldehyde-fixed, paraffin embedded (Mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (KDEL) Polyclonal Antibody, Unconjugated (bs-6940R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (Rat brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (KDEL) Polyclonal Antibody, Unconjugated (bs-6940R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.