

Desmuslin Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP59036**Specification**

Desmuslin Polyclonal Antibody - Product Information

Application	IHC-P, IHC-F, IF, E
Primary Accession	O15061
Reactivity	Rat, Pig, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	172 KDa
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human Desmuslin
Epitope Specificity	501-650/1565
Isotype	IgG
Purity	
affinity purified by Protein A	
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Cytoplasm, cytoskeleton. Cell junction, adherens junction.
SIMILARITY	Belongs to the intermediate filament family.
SUBUNIT	Interacts with GFAP and VIM (By similarity). Isoform 1 interacts with TLN1 and VCL. Isoform 2 interacts with DES and DTNA. Isoform 1 and isoform 2 interact with DMD and UTRN.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Background Descriptions

The protein encoded by this gene is an intermediate filament (IF) family member. IF proteins are cytoskeletal proteins that confer resistance to mechanical stress and are encoded by a dispersed multigene family. This protein has been found to form a linkage between desmin, which is a subunit of the IF network, and the extracellular matrix, and provides an important structural support in muscle. Two alternatively spliced variants encoding different isoforms have been described for this gene. [provided by RefSeq, Jul 2008].

Desmuslin Polyclonal Antibody - Additional Information**Gene ID** 23336**Other Names**

Synemin, Desmuslin, SYNM, DMN, KIAA0353, SYN

Target/Specificity

Isoform 2 is strongly detected in adult heart, fetal skeletal muscles and fetal heart. Isoform 1 is weakly detected in fetal heart and also in fetal skeletal muscle. Isoform 1 and isoform 2 are detected in adult bladder (at protein level). The mRNA is predominantly expressed in heart and muscle with some expression in brain which may be due to tissue-specific isoforms.

Dilution

IHC-P~~N/A
IHC-F~~N/A
IF~~1:50~200
E~~N/A

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Desmuslin Polyclonal Antibody - Protein Information

Name SYNM ([HGNC:24466](#))

Synonyms DMN, KIAA0353, SYN

Function

Type-VI intermediate filament (IF) which plays an important cytoskeletal role within the muscle cell cytoskeleton. It forms heteromeric IFs with desmin and/or vimentin, and via its interaction with cytoskeletal proteins alpha-dystrobrevin, dystrophin, talin-1, utrophin and vinculin, is able to link these heteromeric IFs to adherens-type junctions, such as to the costameres, neuromuscular junctions, and myotendinous junctions within striated muscle cells.

Cellular Location

Cytoplasm, cytoskeleton. Cell junction, adherens junction. Note=There are at least two distinct SYNM subpopulations, one in which SYMN interacts with DES within the Z-lines, and another in which it interacts with both DTNA and DES at the costamere

Tissue Location

Isoform 2 is strongly detected in adult heart, fetal skeletal muscles and fetal heart. Isoform 1 is weakly detected in fetal heart and also in fetal skeletal muscle. Isoform 1 and isoform 2 are detected in adult bladder (at protein level). The mRNA is predominantly expressed in heart and muscle with some expression in brain which may be due to tissue-specific isoforms


Desmuslin Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)

- [Cell Culture](#)

Desmuslin Polyclonal Antibody - Images



Paraformaldehyde-fixed, paraffin embedded (Rat skeletal muscle); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (Desmuslin) Polyclonal Antibody, Unconjugated (bs-8555R) at 1:400 overnight at 4°C, followed by a conjugated Goat Anti-Rabbit IgG antibody (bs-0295G-FITC) for 90 minutes, and DAPI for nuclei staining.