

Anti-Histone Deacetylase 1 (pS421) Antibody

Rabbit polyclonal antibody to Histone Deacetylase 1 (pS421) Catalog # AP59578

Specification

Anti-Histone Deacetylase 1 (pS421) Antibody - Product Information

Application WB
Primary Accession Q13547
Other Accession Q09106

Reactivity Human, Mouse, Rat

Host Rabbit
Clonality Polyclonal
Calculated MW 55103

Anti-Histone Deacetylase 1 (pS421) Antibody - Additional Information

Gene ID 3065

Other Names

RPD3L1; Histone deacetylase 1; HD1

Target/Specificity

Recognizes endogenous levels of Histone Deacetylase 1 (pS421) protein.

Dilution

WB~~WB (1/500 - 1/1000)

Format

Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.

Storage

Store at -20 °C. Stable for 12 months from date of receipt

Anti-Histone Deacetylase 1 (pS421) Antibody - Protein Information

Name HDAC1 {ECO:0000303|PubMed:10846170, ECO:0000312|HGNC:HGNC:4852}

Function

Histone deacetylase that catalyzes the deacetylation of lysine residues on the N-terminal part of the core histones (H2A, H2B, H3 and H4) (PubMed:16762839, PubMed:17704056, PubMed:28497810). Histone deacetylation gives a tag for epigenetic repression and plays an important role in transcriptional regulation, cell cycle progression and developmental events (PubMed:16762839, PubMed:17704056). Histone



deacetylases act via the formation of large multiprotein complexes (PubMed: 16762839, PubMed:17704056). Acts as a component of the histone deacetylase NuRD complex which participates in the remodeling of chromatin (PubMed: 16428440, PubMed:28977666). As part of the SIN3B complex is recruited downstream of the constitutively active genes transcriptional start sites through interaction with histones and mitigates histone acetylation and RNA polymerase II progression within transcribed regions contributing to the regulation of transcription (PubMed:<a $href="http://www.uniprot.org/citations/21041482"\ target="_blank">21041482).\ Also$ functions as a deacetylase for non-histone targets, such as NR1D2, RELA, SP1, SP3, STAT3 and TSHZ3 (PubMed: 12837748, PubMed:16285960, PubMed:16478997, PubMed:17996965, PubMed:19343227). Deacetylates SP proteins, SP1 and SP3, and regulates their function (PubMed:12837748, PubMed:16478997). Component of the BRG1-RB1-HDAC1 complex, which negatively regulates the CREST-mediated transcription in resting neurons (PubMed:19081374). Upon calcium stimulation, HDAC1 is released from the complex and CREBBP is recruited, which facilitates transcriptional activation (PubMed: 19081374). Deacetylates TSHZ3 and regulates its transcriptional repressor activity (PubMed: 19343227). Deacetylates 'Lys-310' in RELA and thereby inhibits the transcriptional activity of NF-kappa-B (PubMed:17000776). Deacetylates NR1D2 and abrogates the effect of KAT5- mediated relieving of NR1D2 transcription repression activity (PubMed: <a href="http://www.uniprot.org/citations/17996965"

target="_blank">17996965). Component of a RCOR/GFI/KDM1A/HDAC complex that suppresses, via histone deacetylase (HDAC) recruitment, a number of genes implicated in multilineage blood cell development (By similarity). Involved in CIART-mediated transcriptional repression of the circadian transcriptional activator: CLOCK-BMAL1 heterodimer (By similarity). Required for the transcriptional repression of circadian target genes, such as PER1, mediated by the large PER complex or CRY1 through histone deacetylation (By similarity). In addition to protein deacetylase activity, also has protein-lysine deacylase activity: acts as a protein decrotonylase and delactylase by mediating decrotonylation ((2E)-butenoyl) and delactylation (lactoyl) of histones, respectively (PubMed:28497810, PubMed:35044827).

Cellular Location Nucleus

Tissue Location

Ubiquitous, with higher levels in heart, pancreas and testis, and lower levels in kidney and brain

Anti-Histone Deacetylase 1 (pS421) Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.


• Western Blot



• Blocking Peptides

- Dot Blot
- Immunohistochemistry
- <u>Immunofluorescence</u>
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

Anti-Histone Deacetylase 1 (pS421) Antibody - Images



Western blot analysis of Histone Deacetylase 1 (pS421) expression in mouse heart (A), rat heart (B), rat testis (C) whole cell lysates.

Anti-Histone Deacetylase 1 (pS421) Antibody - Background

KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human Histone Deacetylase 1. The exact sequence is proprietary.

Anti-Histone Deacetylase 1 (pS421) Antibody - Citations

• ATP50 Hypo-crotonylation Caused by HDAC2 Hyper-Phosphorylation Is a Primary Detrimental Factor for Downregulated Phospholipid Metabolism under Chronic Stress