

Anti-CTIP (pS327) Antibody
Rabbit polyclonal antibody to CTIP (pS327)
Catalog # AP59933

Specification

Anti-CTIP (pS327) Antibody - Product Information

Application	WB
Primary Accession	Q99708
Other Accession	Q80YR6
Reactivity	Human, Mouse, Monkey
Host	Rabbit
Clonality	Polyclonal
Calculated MW	101942

Anti-CTIP (pS327) Antibody - Additional Information

Gene ID 5932

Other Names

CTIP; DNA endonuclease RBBP8; CtBP-interacting protein; CtIP; Retinoblastoma-binding protein 8; RBBP-8; Retinoblastoma-interacting protein and myosin-like; RIM; Sporulation in the absence of SPO11 protein 2 homolog; SAE2

Target/Specificity

KLH-conjugated synthetic peptide encompassing a sequence within the center region of human CTIP (pS327). The exact sequence is proprietary.

Dilution

WB~~WB (1/500 - 1/1000)

Format

Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.

Storage

Store at -20 °C. Stable for 12 months from date of receipt

Anti-CTIP (pS327) Antibody - Protein Information

Name RBBP8

Synonyms CTIP

Function

Endonuclease that cooperates with the MRE11-RAD50-NBN (MRN) complex in DNA-end resection, the first step of double-strand break (DSB) repair through the homologous recombination (HR) pathway (PubMed:17965729, PubMed:19202191)

target="_blank">>19202191, PubMed:>19759395, PubMed:>20064462, PubMed:>23273981, PubMed:>26721387, PubMed:>27814491, PubMed:>27889449, PubMed:>30787182). HR is restricted to S and G2 phases of the cell cycle and preferentially repairs DSBs resulting from replication fork collapse (PubMed:>17965729, PubMed:>19202191, PubMed:>23273981, PubMed:>27814491, PubMed:>27889449, PubMed:>30787182). Key determinant of DSB repair pathway choice, as it commits cells to HR by preventing classical non-homologous end-joining (NHEJ) (PubMed:>19202191). Specifically promotes the endonuclease activity of the MRN complex to clear DNA ends containing protein adducts: recruited to DSBs by NBN following phosphorylation by CDK1, and promotes the endonuclease activity of MRE11 to clear protein-DNA adducts and generate clean double-strand break ends (PubMed:>27814491, PubMed:>27889449, PubMed:>30787182, PubMed:>33836577). Functions downstream of the MRN complex and ATM, promotes ATR activation and its recruitment to DSBs in the S/G2 phase facilitating the generation of ssDNA (PubMed:>16581787, PubMed:>17965729, PubMed:>19759395, PubMed:>20064462). Component of the BRCA1-RBBP8 complex that regulates CHEK1 activation and controls cell cycle G2/M checkpoints on DNA damage (PubMed:>15485915, PubMed:>16818604). During immunoglobulin heavy chain class-switch recombination, promotes microhomology-mediated alternative end joining (A-NHEJ) and plays an essential role in chromosomal translocations (By similarity). Binds preferentially to DNA Y-junctions and to DNA substrates with blocked ends and promotes intermolecular DNA bridging (PubMed:>30601117).

Cellular Location

Nucleus. Chromosome Note=Associates with sites of DNA damage in S/G2 phase (PubMed:10764811, PubMed:25349192). Recruited to DSBs by the MRE11- RAD50-NBN (MRN) complex following phosphorylation by CDK1, which promotes interaction with NBN (PubMed:27814491, PubMed:27889449, PubMed:33836577). Ubiquitinated RBBP8 binds to chromatin following DNA damage (PubMed:16818604).

Tissue Location

Expressed in ER-positive breast cancer lines, but tends to be down-regulated ER-negative cells (at protein level)


Anti-CTIP (pS327) Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)

- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-CTIP (pS327) Antibody - Images



Western blot analysis of CTIP (pS327) expression in U87 (A), PC3 (B), MCF7 (C) whole cell lysates.



Direct ELISA antibody dose-response curve using Anti-CTIP (pS327) Antibody. Antigen (phosphopeptide and non-phosphopeptide) concentration is 5 ug/ml. Goat Anti-Rabbit IgG (H&L) - HRP was used as the secondary antibody, and signal was developed by TMB substrate.

Anti-CTIP (pS327) Antibody - Background

KLH-conjugated synthetic peptide encompassing a sequence within the center region of human CTIP (pS327). The exact sequence is proprietary.