

**Anti-hnRNP G Antibody**  
**Rabbit polyclonal antibody to hnRNP G**  
**Catalog # AP60380****Specification**

---

**Anti-hnRNP G Antibody - Product Information**

Application	WB, IF/IC, IHC
Primary Accession	<a href="#">P38159</a>
Other Accession	<a href="#">Q9WV02</a>
Reactivity	Human, Mouse, Rat, Zebrafish, Pig, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	42332

**Anti-hnRNP G Antibody - Additional Information****Gene ID** 27316**Other Names**

HNRPG; RBMXP1; RNA-binding motif protein, X chromosome; Glycoprotein p43; Heterogeneous nuclear ribonucleoprotein G; hnRNP G

**Target/Specificity**

Recognizes endogenous levels of hnRNP G protein.

**Dilution**

WB~~WB (1/500 - 1/1000), IH (1/100 - 1/200), IF/IC (1/100 - 1/500)

IF/IC~~N/A

IHC~~1:100~500

**Format**

Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.

**Storage**

Store at -20 °C. Stable for 12 months from date of receipt

**Anti-hnRNP G Antibody - Protein Information****Name** RBMX**Synonyms** HNRPG, RBMXP1**Function**

RNA-binding protein that plays several role in the regulation of pre- and post-transcriptional processes. Implicated in tissue- specific regulation of gene transcription and alternative splicing of several pre-mRNAs. Binds to and stimulates transcription from the tumor suppressor TXNIP gene promoter; may thus be involved in tumor suppression. When associated with SAFB, binds to and

stimulates transcription from the SREBF1 promoter. Associates with nascent mRNAs transcribed by RNA polymerase II. Component of the supraspliceosome complex that regulates pre-mRNA alternative splice site selection. Can either activate or suppress exon inclusion; acts additively with TRA2B to promote exon 7 inclusion of the survival motor neuron SMN2. Represses the splicing of MAPT/Tau exon 10. Binds preferentially to single-stranded 5'-CC[A/C]-rich RNA sequence motifs localized in a single-stranded conformation; probably binds RNA as a homodimer. Binds non-specifically to pre-mRNAs. Also plays a role in the cytoplasmic TNFR1 trafficking pathways; promotes both the IL-1-beta-mediated inducible proteolytic cleavage of TNFR1 ectodomains and the release of TNFR1 exosome-like vesicles to the extracellular compartment.

**Cellular Location**

Nucleus Note=Component of ribonucleosomes. Localizes in numerous small granules in the nucleus


**Tissue Location**

Expressed strongly in oral keratinocytes, but only weakly detected in oral squamous cell carcinomas (at protein level)


**Anti-hnRNP G Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**Anti-hnRNP G Antibody - Images**

Western blot analysis of hnRNP G expression in zebrafish (A) whole cell lysates.



Immunohistochemical analysis of hnRNP G staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



Immunofluorescent analysis of hnRNP G staining in Jurkat cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark.

#### **Anti-hnRNP G Antibody - Background**

KLH-conjugated synthetic peptide encompassing a sequence within the N-term region of human hnRNP G. The exact sequence is proprietary.