

Anti-P2Y8 Antibody
Rabbit polyclonal antibody to P2Y8
Catalog # AP60604**Specification**

Anti-P2Y8 Antibody - Product Information

Application	WB, IF/IC
Primary Accession	Q86VZ1
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	40635

Anti-P2Y8 Antibody - Additional Information**Gene ID** 286530**Other Names**

P2Y purinoceptor 8; P2Y8

Target/Specificity

Recognizes endogenous levels of P2Y8 protein.

Dilution

WB~~WB (1/500 - 1/1000), IF/IC (1/100 - 1/500)

IF/IC~~N/A

Format

Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.

Storage

Store at -20 °C. Stable for 12 months from date of receipt

Anti-P2Y8 Antibody - Protein Information**Name** P2RY8 {ECO:0000303|PubMed:30842656, ECO:0000312|HGNC:HGNC:15524}**Function**

G protein-coupled receptor for S-geranylgeranyl-glutathione (GGG), an endogenous metabolite present in lymphoid tissues. Couples the binding of GGG to the activation of GNA13 and downstream repression of AKT activation in lymphocytes defining their positioning and growth within lymphoid organs (PubMed: 25274307, PubMed: 30842656, PubMed: 34088745). In lymphoid follicles, confines B cells and follicular helper T cells in germinal centers (GCs) in response to GGG local gradients established by GGT5 (via GGG catabolism) and ABCC1 (via extracellular transport) with lower concentrations of GGG found in the

follicular dendritic cell network region around which germinal centers are formed (PubMed:25274307, PubMed:30842656, PubMed:34088745). In the bone marrow, also in response to GGG gradients established by GGT5 and ABCC1, it restricts chemotactic transmigration of B cells, T cells and NK cells from blood vessels to the bone marrow parenchyma (PubMed:30842656, PubMed:34088745). Contributes to GNA13-dependent pathway that suppresses GC B cell growth (PubMed:25274307).

Cellular Location

Cell membrane; Multi-pass membrane protein

Tissue Location


Barely detectable in normal blood leukocytes. Weaker expression was seen in heart, kidney and lung. Not detected in brain (PubMed:11004484, PubMed:15466006). Expressed in B cells and follicular helper T cells in germinal centers (at protein level) (PubMed:30842656).

Anti-P2Y8 Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-P2Y8 Antibody - Images



Western blot analysis of P2Y8 expression in Hela (A), mouse lung (B), mouse heart (C), mouse kidney (D), rat heart (E), rat kidney (F) whole cell lysates.



Immunofluorescent analysis of P2Y8 staining in HuvEc cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).

Anti-P2Y8 Antibody - Background

KLH-conjugated synthetic peptide encompassing a sequence within the center region of human P2Y8. The exact sequence is proprietary.