

**Anti-Parkin Antibody**  
**Rabbit polyclonal antibody to Parkin**  
**Catalog # AP60605**

## Specification

## **Anti-Parkin Antibody - Product Information**

Application	WB, IF/IC, IHC
Primary Accession	<a href="#">060260</a>
Reactivity	Human, Mouse, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	51641

## **Anti-Parkin Antibody - Additional Information**

Gene ID 5071

## Other Names

PRKN; E3 ubiquitin-protein ligase parkin; Parkinson juvenile disease protein 2; Parkinson disease protein 2

## Target/Specificity

Recognizes endogenous levels of Parkin protein.

## Dilution

WB~~WB (1/500 - 1/1000), IH (1/100 - 1/200), IF/IC (1/100 - 1/500)  
IF/IC~~N/A  
IHC~~1:100~500

## Format

Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.

## Storage

Store at -20 °C. Stable for 12 months from date of receipt

## Anti-Parkin Antibody - Protein Information

Name PRKN ([HGNC:8607](#))

## Synonyms PARK2

## Function

href="http://www.uniprot.org/citations/12150907" target="\_blank">>12150907</a>, PubMed:<a href="http://www.uniprot.org/citations/12628165" target="\_blank">>12628165</a>, PubMed:<a href="http://www.uniprot.org/citations/15105460" target="\_blank">>15105460</a>, PubMed:<a href="http://www.uniprot.org/citations/16135753" target="\_blank">>16135753</a>, PubMed:<a href="http://www.uniprot.org/citations/21376232" target="\_blank">>21376232</a>, PubMed:<a href="http://www.uniprot.org/citations/21532592" target="\_blank">>21532592</a>, PubMed:<a href="http://www.uniprot.org/citations/22396657" target="\_blank">>22396657</a>, PubMed:<a href="http://www.uniprot.org/citations/23620051" target="\_blank">>23620051</a>, PubMed:<a href="http://www.uniprot.org/citations/23754282" target="\_blank">>23754282</a>, PubMed:<a href="http://www.uniprot.org/citations/24660806" target="\_blank">>24660806</a>, PubMed:<a href="http://www.uniprot.org/citations/24751536" target="\_blank">>24751536</a>, PubMed:<a href="http://www.uniprot.org/citations/29311685" target="\_blank">>29311685</a>, PubMed:<a href="http://www.uniprot.org/citations/32047033" target="\_blank">>32047033</a>). Substrates include SYT11 and VDAC1 (PubMed:<a href="http://www.uniprot.org/citations/29311685" target="\_blank">>29311685</a>, PubMed:<a href="http://www.uniprot.org/citations/32047033" target="\_blank">>32047033</a>). Other substrates are BCL2, CCNE1, GPR37, RHOT1/MIRO1, MFN1, MFN2, STUB1, SNCAIP, SEPTIN5, TOMM20, USP30, ZNF746, MIRO1 and AIMP2 (PubMed:<a href="http://www.uniprot.org/citations/10888878" target="\_blank">>10888878</a>, PubMed:<a href="http://www.uniprot.org/citations/10973942" target="\_blank">>10973942</a>, PubMed:<a href="http://www.uniprot.org/citations/11431533" target="\_blank">>11431533</a>, PubMed:<a href="http://www.uniprot.org/citations/12150907" target="\_blank">>12150907</a>, PubMed:<a href="http://www.uniprot.org/citations/12628165" target="\_blank">>12628165</a>, PubMed:<a href="http://www.uniprot.org/citations/15105460" target="\_blank">>15105460</a>, PubMed:<a href="http://www.uniprot.org/citations/16135753" target="\_blank">>16135753</a>, PubMed:<a href="http://www.uniprot.org/citations/21376232" target="\_blank">>21376232</a>, PubMed:<a href="http://www.uniprot.org/citations/21532592" target="\_blank">>21532592</a>, PubMed:<a href="http://www.uniprot.org/citations/22396657" target="\_blank">>22396657</a>, PubMed:<a href="http://www.uniprot.org/citations/23620051" target="\_blank">>23620051</a>, PubMed:<a href="http://www.uniprot.org/citations/23754282" target="\_blank">>23754282</a>, PubMed:<a href="http://www.uniprot.org/citations/24660806" target="\_blank">>24660806</a>, PubMed:<a href="http://www.uniprot.org/citations/24751536" target="\_blank">>24751536</a>). Mediates monoubiquitination as well as 'Lys-6', 'Lys-11', 'Lys-48'-linked and 'Lys-63'-linked polyubiquitination of substrates depending on the context (PubMed:<a href="http://www.uniprot.org/citations/19229105" target="\_blank">>19229105</a>, PubMed:<a href="http://www.uniprot.org/citations/20889974" target="\_blank">>20889974</a>, PubMed:<a href="http://www.uniprot.org/citations/25474007" target="\_blank">>25474007</a>, PubMed:<a href="http://www.uniprot.org/citations/25621951" target="\_blank">>25621951</a>, PubMed:<a href="http://www.uniprot.org/citations/32047033" target="\_blank">>32047033</a>). Participates in the removal and/or detoxification of abnormally folded or damaged protein by mediating 'Lys-63'-linked polyubiquitination of misfolded proteins such as PARK7: 'Lys-63'-linked polyubiquitinated misfolded proteins are then recognized by HDAC6, leading to their recruitment to aggresomes, followed by degradation (PubMed:<a href="http://www.uniprot.org/citations/17846173" target="\_blank">>17846173</a>, PubMed:<a href="http://www.uniprot.org/citations/19229105" target="\_blank">>19229105</a>). Mediates 'Lys-63'-linked polyubiquitination of a 22 kDa O-linked glycosylated isoform of SNCAIP, possibly playing a role in Lewy-body formation (PubMed:<a href="http://www.uniprot.org/citations/11431533" target="\_blank">>11431533</a>, PubMed:<a href="http://www.uniprot.org/citations/11590439" target="\_blank">>11590439</a>, PubMed:<a href="http://www.uniprot.org/citations/15105460" target="\_blank">>15105460</a>, PubMed:<a href="http://www.uniprot.org/citations/15728840" target="\_blank">>15728840</a>, PubMed:<a href="http://www.uniprot.org/citations/19229105" target="\_blank">>19229105</a>). Mediates monoubiquitination of BCL2, thereby acting as a positive regulator of autophagy (PubMed:<a href="http://www.uniprot.org/citations/20889974" target="\_blank">>20889974</a>). Protects against mitochondrial dysfunction during cellular stress, by acting downstream of PINK1 to coordinate mitochondrial quality control mechanisms that remove and replace dysfunctional mitochondrial components (PubMed:<a href="http://www.uniprot.org/citations/11439185" target="\_blank">>11439185</a>, PubMed:<a href="http://www.uniprot.org/citations/18957282" target="\_blank">>11439185</a>),

target="\_blank">>18957282</a>, PubMed:<a href="http://www.uniprot.org/citations/19029340" target="\_blank">19029340</a>, PubMed:<a href="http://www.uniprot.org/citations/19966284" target="\_blank">19966284</a>, PubMed:<a href="http://www.uniprot.org/citations/21376232" target="\_blank">21376232</a>, PubMed:<a href="http://www.uniprot.org/citations/22082830" target="\_blank">22082830</a>, PubMed:<a href="http://www.uniprot.org/citations/22396657" target="\_blank">22396657</a>, PubMed:<a href="http://www.uniprot.org/citations/23620051" target="\_blank">23620051</a>, PubMed:<a href="http://www.uniprot.org/citations/23933751" target="\_blank">23933751</a>, PubMed:<a href="http://www.uniprot.org/citations/24660806" target="\_blank">24660806</a>, PubMed:<a href="http://www.uniprot.org/citations/24784582" target="\_blank">24784582</a>, PubMed:<a href="http://www.uniprot.org/citations/24896179" target="\_blank">24896179</a>, PubMed:<a href="http://www.uniprot.org/citations/25474007" target="\_blank">25474007</a>, PubMed:<a href="http://www.uniprot.org/citations/25527291" target="\_blank">25527291</a>, PubMed:<a href="http://www.uniprot.org/citations/32047033" target="\_blank">32047033</a>). Depending on the severity of mitochondrial damage and/or dysfunction, activity ranges from preventing apoptosis and stimulating mitochondrial biogenesis to regulating mitochondrial dynamics and eliminating severely damaged mitochondria via mitophagy (PubMed:<a href="http://www.uniprot.org/citations/11439185" target="\_blank">11439185</a>, PubMed:<a href="http://www.uniprot.org/citations/19029340" target="\_blank">19029340</a>, PubMed:<a href="http://www.uniprot.org/citations/19801972" target="\_blank">19801972</a>, PubMed:<a href="http://www.uniprot.org/citations/19966284" target="\_blank">19966284</a>, PubMed:<a href="http://www.uniprot.org/citations/21376232" target="\_blank">21376232</a>, PubMed:<a href="http://www.uniprot.org/citations/22082830" target="\_blank">22082830</a>, PubMed:<a href="http://www.uniprot.org/citations/22396657" target="\_blank">22396657</a>, PubMed:<a href="http://www.uniprot.org/citations/23620051" target="\_blank">23620051</a>, PubMed:<a href="http://www.uniprot.org/citations/23685073" target="\_blank">23685073</a>, PubMed:<a href="http://www.uniprot.org/citations/23933751" target="\_blank">23933751</a>, PubMed:<a href="http://www.uniprot.org/citations/24896179" target="\_blank">24896179</a>, PubMed:<a href="http://www.uniprot.org/citations/25527291" target="\_blank">25527291</a>, PubMed:<a href="http://www.uniprot.org/citations/32047033" target="\_blank">32047033</a>, PubMed:<a href="http://www.uniprot.org/citations/33499712" target="\_blank">33499712</a>). Activation and recruitment onto the outer membrane of damaged/dysfunctional mitochondria (OMM) requires PINK1-mediated phosphorylation of both PRKN and ubiquitin (PubMed:<a href="http://www.uniprot.org/citations/24660806" target="\_blank">24660806</a>, PubMed:<a href="http://www.uniprot.org/citations/24784582" target="\_blank">24784582</a>, PubMed:<a href="http://www.uniprot.org/citations/25474007" target="\_blank">25474007</a>, PubMed:<a href="http://www.uniprot.org/citations/25527291" target="\_blank">25527291</a>). After mitochondrial damage, functions with PINK1 to mediate the decision between mitophagy or preventing apoptosis by inducing either the poly- or monoubiquitination of VDAC1, respectively; polyubiquitination of VDAC1 promotes mitophagy, while monoubiquitination of VDAC1 decreases mitochondrial calcium influx which ultimately inhibits apoptosis (PubMed:<a href="http://www.uniprot.org/citations/27534820" target="\_blank">27534820</a>, PubMed:<a href="http://www.uniprot.org/citations/32047033" target="\_blank">32047033</a>). When cellular stress results in irreversible mitochondrial damage, promotes the autophagic degradation of dysfunctional depolarized mitochondria (mitophagy) by promoting the ubiquitination of mitochondrial proteins such as TOMM20, RHOT1/MIRO1, MFN1 and USP30 (PubMed:<a href="http://www.uniprot.org/citations/19029340" target="\_blank">19029340</a>, PubMed:<a href="http://www.uniprot.org/citations/19966284" target="\_blank">19966284</a>, PubMed:<a href="http://www.uniprot.org/citations/21753002" target="\_blank">21753002</a>, PubMed:<a href="http://www.uniprot.org/citations/22396657" target="\_blank">22396657</a>, PubMed:<a href="http://www.uniprot.org/citations/23620051" target="\_blank">23620051</a>, PubMed:<a href="http://www.uniprot.org/citations/23685073" target="\_blank">23685073</a>, PubMed:<a href="http://www.uniprot.org/citations/23933751" target="\_blank">23933751</a>, PubMed:<a href="http://www.uniprot.org/citations/24896179" target="\_blank">24896179</a>, PubMed:<a href="http://www.uniprot.org/citations/25527291" target="\_blank">25527291</a>). Preferentially assembles 'Lys-6'-, 'Lys-11'- and 'Lys-63'-linked polyubiquitin chains, leading to mitophagy (PubMed:<a href="http://www.uniprot.org/citations/25621951" target="\_blank">25621951</a>, PubMed:<a href="http://www.uniprot.org/citations/32047033" target="\_blank">32047033</a>).

The PINK1-PRKN pathway also promotes fission of damaged mitochondria by PINK1-mediated phosphorylation which promotes the PRKN-dependent degradation of mitochondrial proteins involved in fission such as MFN2 (PubMed:<a href="http://www.uniprot.org/citations/23620051" target="\_blank">23620051</a>). This prevents the refusion of unhealthy mitochondria with the mitochondrial network or initiates mitochondrial fragmentation facilitating their later engulfment by autophagosomes (PubMed:<a href="http://www.uniprot.org/citations/23620051" target="\_blank">23620051</a>). Regulates motility of damaged mitochondria via the ubiquitination and subsequent degradation of MIRO1 and MIRO2; in motor neurons, this likely inhibits mitochondrial intracellular anterograde transport along the axons which probably increases the chance of the mitochondria undergoing mitophagy in the soma (PubMed:<a href="http://www.uniprot.org/citations/22396657" target="\_blank">22396657</a>). Involved in mitochondrial biogenesis via the 'Lys-48'-linked polyubiquitination of transcriptional repressor ZNF746/PARIS which leads to its subsequent proteasomal degradation and allows activation of the transcription factor PPARC1A (PubMed:<a href="http://www.uniprot.org/citations/21376232" target="\_blank">21376232</a>). Limits the production of reactive oxygen species (ROS) (PubMed:<a href="http://www.uniprot.org/citations/18541373" target="\_blank">18541373</a>). Regulates cyclin-E during neuronal apoptosis (PubMed:<a href="http://www.uniprot.org/citations/12628165" target="\_blank">12628165</a>). In collaboration with CHPF isoform 2, may enhance cell viability and protect cells from oxidative stress (PubMed:<a href="http://www.uniprot.org/citations/22082830" target="\_blank">22082830</a>). Independently of its ubiquitin ligase activity, protects from apoptosis by the transcriptional repression of p53/TP53 (PubMed:<a href="http://www.uniprot.org/citations/19801972" target="\_blank">19801972</a>). May protect neurons against alpha synuclein toxicity, proteasomal dysfunction, GPR37 accumulation, and kainate-induced excitotoxicity (PubMed:<a href="http://www.uniprot.org/citations/11439185" target="\_blank">11439185</a>). May play a role in controlling neurotransmitter trafficking at the presynaptic terminal and in calcium-dependent exocytosis. May represent a tumor suppressor gene (PubMed:<a href="http://www.uniprot.org/citations/12719539" target="\_blank">12719539</a>).

### Cellular Location

Cytoplasm, cytosol. Nucleus. Endoplasmic reticulum. Mitochondrion. Mitochondrion outer membrane {ECO:0000250|UniProtKB:Q9WVS6}. Cell projection, neuron projection. Postsynaptic density {ECO:0000250|UniProtKB:Q9WVS6}. Presynapse {ECO:0000250|UniProtKB:Q9WVS6}. Note=Mainly localizes in the cytosol (PubMed:19029340, PubMed:19229105). Co-localizes with SYT11 in neutrites (PubMed:12925569). Co-localizes with SNCAIP in brainstem Lewy bodies (PubMed:10319893, PubMed:11431533). Translocates to dysfunctional mitochondria that have lost the mitochondrial membrane potential; recruitment to mitochondria is PINK1-dependent (PubMed:18957282, PubMed:19966284, PubMed:23620051, PubMed:24898855) Mitochondrial localization also gradually increases with cellular growth (PubMed:22082830).

### Tissue Location


Highly expressed in the brain including the substantia nigra (PubMed:19501131, PubMed:9560156). Expressed in heart, testis and skeletal muscle (PubMed:9560156). Expression is down- regulated or absent in tumor biopsies, and absent in the brain of PARK2 patients (PubMed:12719539, PubMed:14614460). Overexpression protects dopamine neurons from kainate-mediated apoptosis (PubMed:12628165) Found in serum (at protein level) (PubMed:19501131)

### Anti-Parkin Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)


- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**Anti-Parkin Antibody - Images**

Western blot analysis of Parkin expression in mouse kidney (A), mouse heart (B) whole cell lysates.



Immunohistochemical analysis of Parkin staining in human brain formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



Immunofluorescent analysis of Parkin staining in U87MG cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).

#### **Anti-Parkin Antibody - Background**

KLH-conjugated synthetic peptide encompassing a sequence within the center region of human Parkin. The exact sequence is proprietary.