

Anti-FGFR4 Antibody
Rabbit polyclonal antibody to FGFR4
Catalog # AP60670**Specification**

Anti-FGFR4 Antibody - Product Information

Application	WB
Primary Accession	P22455
Other Accession	Q03142
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Calculated MW	87954

Anti-FGFR4 Antibody - Additional Information**Gene ID** 2264**Other Names**

JTK2; TKF; Fibroblast growth factor receptor 4; FGFR-4; CD334

Target/Specificity

Recognizes endogenous levels of FGFR4 protein.

Dilution

WB~~WB (1/500 - 1/1000)

Format

Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.

Storage

Store at -20 °C. Stable for 12 months from date of receipt

Anti-FGFR4 Antibody - Protein Information**Name** FGFR4**Synonyms** JTK2, TKF**Function**

Tyrosine-protein kinase that acts as a cell-surface receptor for fibroblast growth factors and plays a role in the regulation of cell proliferation, differentiation and migration, and in regulation of lipid metabolism, bile acid biosynthesis, glucose uptake, vitamin D metabolism and phosphate homeostasis. Required for normal down- regulation of the expression of CYP7A1, the rate-limiting enzyme in bile acid synthesis, in response to FGF19. Phosphorylates PLCG1 and FRS2. Ligand binding leads to the activation of several signaling cascades. Activation of PLCG1 leads to the production of the cellular signaling molecules diacylglycerol and inositol 1,4,5-trisphosphate.

Phosphorylation of FRS2 triggers recruitment of GRB2, GAB1, PIK3R1 and SOS1, and mediates activation of RAS, MAPK1/ERK2, MAPK3/ERK1 and the MAP kinase signaling pathway, as well as of the AKT1 signaling pathway. Promotes SRC-dependent phosphorylation of the matrix protease MMP14 and its lysosomal degradation. FGFR4 signaling is down-regulated by receptor internalization and degradation; MMP14 promotes internalization and degradation of FGFR4. Mutations that lead to constitutive kinase activation or impair normal FGFR4 inactivation lead to aberrant signaling.

Cellular Location

Cell membrane; Single-pass type I membrane protein. Endosome. Endoplasmic reticulum.

Note=Internalized from the cell membrane to recycling endosomes, and from there back to the cell membrane


Tissue Location

Expressed in gastrointestinal epithelial cells, pancreas, and gastric and pancreatic cancer cell lines

Anti-FGFR4 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-FGFR4 Antibody - Images

Western blot analysis of FGFR4 expression in HepG2 (A), HEK293T (B), AML12 (C) whole cell lysates.

Anti-FGFR4 Antibody - Background

KLH-conjugated synthetic peptide encompassing a sequence within the center region of human FGFR4. The exact sequence is proprietary.