

α -tubulin (Acetyl Lys40) Monoclonal Antibody(4A8)
Catalog # AP63647**Specification**

 α -tubulin (Acetyl Lys40) Monoclonal Antibody(4A8) - Product Information

Application	WB, IHC-P, IF
Primary Accession	P68363
Reactivity	Human, Rat, Mouse
Host	Mouse
Clonality	Monoclonal

 α -tubulin (Acetyl Lys40) Monoclonal Antibody(4A8) - Additional Information**Gene ID** 10376**Other Names**

Tubulin alpha-1B chain (Alpha-tubulin ubiquitous) (Tubulin K-alpha-1) (Tubulin alpha-ubiquitous chain)

Dilution

WB~~WB 1:1000-2000, IHC 1:50-100 IF 1:200

IHC-P~~N/A

IF~~1:50~200

Format

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

Storage Conditions

-20°C

 α -tubulin (Acetyl Lys40) Monoclonal Antibody(4A8) - Protein Information**Name** TUBA1B**Function**

Tubulin is the major constituent of microtubules, a cylinder consisting of laterally associated linear protofilaments composed of alpha- and beta-tubulin heterodimers (PubMed:34996871). Microtubules grow by the addition of GTP-tubulin dimers to the microtubule end, where a stabilizing cap forms (PubMed:34996871). Below the cap, tubulin dimers are in GDP-bound state, owing to GTPase activity of alpha- tubulin (PubMed:34996871).

Cellular Location




Cytoplasm, cytoskeleton






α -tubulin (Acetyl Lys40) Monoclonal Antibody(4A8) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

α -tubulin (Acetyl Lys40) Monoclonal Antibody(4A8) - Images







α -tubulin (Acetyl Lys40) Monoclonal Antibody(4A8) - Background

Tubulin is the major constituent of microtubules. It binds two moles of GTP, one at an exchangeable site on the beta chain and one at a non-exchangeable site on the alpha chain.

α -tubulin (Acetyl Lys40) Monoclonal Antibody(4A8) - Citations

- [DDHD1, but Not DDHD2, Suppresses Neurite Outgrowth in SH-SY5Y and PC12 Cells by Regulating Protein Transport From Recycling Endosomes](#)