

Histone H2A.X (phospho Ser139) Polyclonal Antibody
Catalog # AP67060**Specification**

Histone H2A.X (phospho Ser139) Polyclonal Antibody - Product Information

Application	WB, IHC-P
Primary Accession	P16104
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal

Histone H2A.X (phospho Ser139) Polyclonal Antibody - Additional Information**Gene ID** 3014**Other Names**

H2AFX; H2AX; Histone H2A.x; H2a/x

Dilution

WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/10000. Not yet tested in other applications.

IHC-P~~N/A

Format

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

Storage Conditions

-20°C

Histone H2A.X (phospho Ser139) Polyclonal Antibody - Protein Information**Name** H2AX ([HGNC:4739](#))**Function**

Variant histone H2A which replaces conventional H2A in a subset of nucleosomes. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post- translational modifications of histones, also called histone code, and nucleosome remodeling. Required for checkpoint-mediated arrest of cell cycle progression in response to low doses of ionizing radiation and for efficient repair of DNA double strand breaks (DSBs) specifically when modified by C-terminal phosphorylation.

Cellular Location




Nucleus. Chromosome





Histone H2A.X (phospho Ser139) Polyclonal Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Histone H2A.X (phospho Ser139) Polyclonal Antibody - Images







Histone H2A.X (phospho Ser139) Polyclonal Antibody - Background

Variant histone H2A which replaces conventional H2A in a subset of nucleosomes. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of histones, also called histone code, and nucleosome remodeling. Required for checkpoint-mediated arrest of cell cycle progression in response to low doses of ionizing radiation and for efficient repair of DNA double strand breaks (DSBs) specifically when modified by C- terminal phosphorylation.