

TOMM70A Antibody (C-term)

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP6745b

Specification

TOMM70A Antibody (C-term) - Product Information

Application FC, IHC-P, WB,E

Primary Accession <u>094826</u>

Other Accession <u>Q75Q39</u>, <u>Q9CZW5</u>

Reactivity
Predicted
Host
Clonality
Isotype
Antigen Region

Human
Mouse, Rat
Rabbit
Polyclonal
Rabbit IgG
579-608

TOMM70A Antibody (C-term) - Additional Information

Gene ID 9868

Other Names

Mitochondrial import receptor subunit TOM70, Mitochondrial precursor proteins import receptor, Translocase of outer membrane 70 kDa subunit, TOMM70A, KIAA0719, TOM70

Target/Specificity

This TOMM70A antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 579-608 amino acids from the C-terminal region of human TOMM70A.

Dilution

FC~~1:10~50 IHC-P~~1:50~100 WB~~1:1000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

TOMM70A Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

TOMM70A Antibody (C-term) - Protein Information





Name TOMM70 (<u>HGNC:11985</u>)

Function Acts as a receptor of the preprotein translocase complex of the outer mitochondrial membrane (TOM complex) (PubMed:12526792). Recognizes and mediates the translocation of mitochondrial preproteins from the cytosol into the mitochondria in a chaperone dependent manner (PubMed:12526792, PubMed:35025629). Mediates TBK1 and IRF3 activation induced by MAVS in response to Sendai virus infection and promotes host antiviral responses during virus infection (PubMed:20628368, PubMed:25609812, PubMed:32728199). Upon Sendai virus infection, recruits HSP90AA1:IRF3:BAX in mitochondrion and the complex induces apoptosis (PubMed:25609812).

Cellular Location


Mitochondrion outer membrane; Single-pass membrane protein

TOMM70A Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

TOMM70A Antibody (C-term) - Images




Western blot analysis of TOMM70A antibody (C-term) (Cat. #AP6745b) in HL60 cell line lysates (35ug/lane). TOMM70A (arrow) was detected using the purified Pab.





Formalin-fixed and paraffin-embedded human brain tissue reacted with TOMM70A Antibody (C-term), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.



Flow cytometric analysis of hela cells using TOMM70A Antibody (C-term)(bottom histogram) compared to a negative control cell (top histogram)FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

TOMM70A Antibody (C-term) - Background

The translocase of outer mitochondrial membrane (TOM) complex is a multisubunit complex involved in the recognition, unfolding, and translocation of preproteins into the mitochondria.

TOMM70A Antibody (C-term) - References

Blesa, J.R., Gene 427 (1-2), 58-64 (2008) Chou, C.H., Mol. Biol. Cell 17 (9), 3952-3963 (2006) Edmonson, A.M., Cell Commun. Adhes. 9 (1), 15-27 (2002)

TOMM70A Antibody (C-term) - Citations

- Endothelial Mitochondrial Preprotein Translocase Tomm7-Rac1 Signaling Axis Dominates Cerebrovascular Network Homeostasis.
- Proteomic analysis of mitochondria in respiratory epithelial cells infected with human respiratory syncytial virus and functional implications for virus and cell biology.