

Ku70 Antibody (Center)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP6775c**Specification**

Ku70 Antibody (Center) - Product Information

Application	FC, IHC-P, WB,E
Primary Accession	P12956
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	69843
Antigen Region	432-461

Ku70 Antibody (Center) - Additional Information**Gene ID** 2547**Other Names**

X-ray repair cross-complementing protein 6, 364-, 4299-, 5'-deoxyribose-5-phosphate lyase Ku70, 5'-dRP lyase Ku70, 70 kDa subunit of Ku antigen, ATP-dependent DNA helicase 2 subunit 1, ATP-dependent DNA helicase II 70 kDa subunit, CTC box-binding factor 75 kDa subunit, CTC75, CTCBF, DNA repair protein XRCC6, Lupus Ku autoantigen protein p70, Ku70, Thyroid-lupus autoantigen, TLAA, X-ray repair complementing defective repair in Chinese hamster cells 6, XRCC6, G22P1

Target/Specificity

This Ku70 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 432-461 amino acids from the Central region of human Ku70.

Dilution

FC~~1:10~50

IHC-P~~1:10~50

WB~~1:1000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Ku70 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

Ku70 Antibody (Center) - Protein Information

Name XRCC6

Synonyms G22P1

Function Single-stranded DNA-dependent ATP-dependent helicase that plays a key role in DNA non-homologous end joining (NHEJ) by recruiting DNA-PK to DNA (PubMed:[11493912](#), PubMed:[12145306](#), PubMed:[20493174](#), PubMed:[2466842](#), PubMed:[7957065](#), PubMed:[8621488](#), PubMed:[9742108](#)). Required for double-strand break repair and V(D)J recombination (PubMed:[11493912](#), PubMed:[12145306](#), PubMed:[20493174](#), PubMed:[2466842](#), PubMed:[7957065](#), PubMed:[8621488](#), PubMed:[9742108](#)). Also has a role in chromosome translocation (PubMed:[11493912](#), PubMed:[12145306](#), PubMed:[20493174](#), PubMed:[2466842](#), PubMed:[7957065](#), PubMed:[8621488](#), PubMed:[9742108](#)). Has a role in chromosome translocation (PubMed:[11493912](#), PubMed:[12145306](#), PubMed:[20493174](#), PubMed:[2466842](#), PubMed:[7957065](#), PubMed:[8621488](#), PubMed:[9742108](#)). The DNA helicase II complex binds preferentially to fork-like ends of double-stranded DNA in a cell cycle-dependent manner (PubMed:[11493912](#), PubMed:[12145306](#), PubMed:[20493174](#), PubMed:[2466842](#), PubMed:[7957065](#), PubMed:[8621488](#), PubMed:[9742108](#)). It works in the 3'-5' direction (PubMed:[11493912](#), PubMed:[12145306](#), PubMed:[20493174](#), PubMed:[2466842](#), PubMed:[7957065](#), PubMed:[8621488](#), PubMed:[9742108](#)). During NHEJ, the XRCC5-XRCC6 dimer performs the recognition step: it recognizes and binds to the broken ends of the DNA and protects them from further resection (PubMed:[11493912](#), PubMed:[12145306](#), PubMed:[20493174](#), PubMed:[2466842](#), PubMed:[7957065](#), PubMed:[8621488](#), PubMed:[9742108](#)). Binding to DNA may be mediated by XRCC6 (PubMed:[11493912](#), PubMed:[12145306](#), PubMed:[20493174](#), PubMed:[2466842](#), PubMed:[7957065](#), PubMed:[8621488](#), PubMed:[9742108](#)). The XRCC5-XRCC6 dimer acts as a regulatory subunit of the DNA-dependent protein kinase complex DNA-PK by increasing the affinity of the catalytic subunit PRKDC to DNA by 100-fold (PubMed:[11493912](#), PubMed:[12145306](#), PubMed:[20493174](#), PubMed:[2466842](#), PubMed:[7957065](#), PubMed:[8621488](#), PubMed:[9742108](#)). The XRCC5-XRCC6 dimer is probably involved in stabilizing broken DNA ends and bringing them together (PubMed:[11493912](#), PubMed:[12145306](#), PubMed:[20493174](#), PubMed:[2466842](#), PubMed:[7957065](#), PubMed:[8621488](#), PubMed:[9742108](#)). The assembly of the DNA-PK complex to DNA ends is required for the NHEJ ligation step (PubMed:[11493912](#), PubMed:[12145306](#), PubMed:[20493174](#), PubMed:[2466842](#), PubMed:[7957065](#), PubMed:[8621488](#), PubMed:[9742108](#)). Probably also acts as a 5'-deoxyribose-5-phosphate lyase (5'-dRP lyase), by catalyzing the beta-elimination of the 5' deoxyribose-5-phosphate at an abasic site near double-strand breaks (PubMed:[20383123](#)). 5'-dRP lyase activity allows to 'clean' the termini of abasic sites, a class of nucleotide damage commonly associated with strand breaks, before such broken ends can be joined (PubMed:[20383123](#)). The XRCC5-XRCC6 dimer together with APEX1 acts as a negative regulator of transcription (PubMed:[8621488](#)). In association with NAA15, the XRCC5-XRCC6 dimer binds to the osteocalcin promoter and activates osteocalcin expression (PubMed:[12145306](#)). Plays a role in the regulation of DNA virus-mediated innate immune response by assembling into the HDP-RNP complex, a complex that serves as a platform for IRF3 phosphorylation and subsequent innate immune response activation through the cGAS-STING pathway (PubMed:[28712728](#)). Negatively regulates apoptosis by interacting with BAX and sequestering it from the mitochondria (PubMed:[15023334](#)). Might have deubiquitination activity, acting on BAX (PubMed:[18362350](#)).

Cellular Location

Nucleus. Chromosome. Cytoplasm. Note=When trimethylated, localizes in the cytoplasm.


Ku70 Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)

- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


Ku70 Antibody (Center) - Images



Western blot analysis of Ku70 (arrow) using rabbit polyclonal Ku70 Antibody (Center) (Cat. #AP6775c). 293 cell lysates (2 ug/lane) either nontransfected (Lane 1) or transiently transfected with the Ku70 gene (Lane 2).



Formalin-fixed and paraffin-embedded mouse spleen tissue reacted with Ku70 Antibody (Center), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.



Ku70 Antibody (Center) (Cat. #AP6775c) flow cytometric analysis of 293 cells (right histogram) compared to a negative control cell (left histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

Ku70 Antibody (Center) - Background

Ku70 is a single stranded DNA-dependent ATP-dependent helicase. It has a role in chromosome translocation. The DNA helicase II complex binds preferentially to fork-like ends of double-stranded DNA in a cell cycle-dependent manner. It works in the 3'-5' direction. Binding to DNA may be mediated by p70. It is involved in DNA nonhomologous end joining (NHEJ) required for double-strand break repair and V(D)J recombination. The Ku p70/p86 dimer acts as regulatory subunit of the DNA-dependent protein kinase complex DNA-PK by increasing the affinity of the catalytic subunit PRKDC to DNA by 100-fold. The Ku p70/p86 dimer is probably involved in stabilizing broken DNA ends and bringing them together.

Ku70 Antibody (Center) - References

Beskow, C., et al., Br. J. Cancer 101 (5), 816-821 (2009) Tseng, R.C., et al., Cancer 115 (13), 2939-2948 (2009)

Ku70 Antibody (Center) - Citations

- [JmjC domain-containing protein 8 \(JMJD8\) represses Ku70/Ku80 expression via attenuating AKT/NF-κB/COX-2 signaling.](#)