

SNRPD2 Antibody (N-term)

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP6841a

Specification

SNRPD2 Antibody (N-term) - Product Information

Application Primary Accession Other Accession Reactivity Predicted Host Clonality Isotype Calculated MW Antigen Region FC, IHC-P, WB,E <u>P62316</u> <u>O29329</u>, <u>P62317</u>, <u>O3SZF8</u> Human Bovine, Mouse, Pig Rabbit Polyclonal Rabbit IgG 13527 1-30

SNRPD2 Antibody (N-term) - Additional Information

Gene ID 6633

Other Names Small nuclear ribonucleoprotein Sm D2, Sm-D2, snRNP core protein D2, SNRPD2, SNRPD1

Target/Specificity

This SNRPD2 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 1-30 amino acids from the N-terminal region of human SNRPD2.

Dilution FC~~1:10~50 IHC-P~~1:50~100 WB~~1:1000 E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

SNRPD2 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

SNRPD2 Antibody (N-term) - Protein Information



Name SNRPD2

Synonyms SNRPD1

Function Plays a role in pre-mRNA splicing as a core component of the spliceosomal U1, U2, U4 and U5 small nuclear ribonucleoproteins (snRNPs), the building blocks of the spliceosome (PubMed:<u>11991638</u>, PubMed:<u>18984161</u>, PubMed:<u>19325628</u>, PubMed:<u>23333303</u>, PubMed:<u>25555158</u>, PubMed:<u>26912367</u>, PubMed:<u>28076346</u>, PubMed:<u>28502770</u>, PubMed:<u>28781166</u>, PubMed:<u>32494006</u>). Component of both the pre-catalytic spliceosome B complex and activated spliceosome C complexes (PubMed:<u>11991638</u>, PubMed:<u>28076346</u>, PubMed:<u>28502770</u>, PubMed:<u>28502770</u>, PubMed:<u>28781166</u>). As a component of the minor spliceosome, involved in the splicing of U12-type introns in pre-mRNAs (PubMed:<u>15146077</u>).

Cellular Location

Cytoplasm, cytosol. Nucleus. Note=SMN-mediated assembly into core snRNPs occurs in the cytosol before SMN-mediated transport to the nucleus to be included in spliceosomes.

SNRPD2 Antibody (N-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

SNRPD2 Antibody (N-term) - Images


не 55	ela
28	
17	•4
11	

Western blot analysis of SNRPD2 Antibody (N-term) (Cat. #AP6841a) in Hela cell line lysates (35ug/lane). SNRPD2 (arrow) was detected using the purified Pab.





Formalin-fixed and paraffin-embedded human lung carcinoma reacted with SNRPD2 Antibody (N-term), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.



SNRPD2 Antibody (N-term) (Cat. #AP6841a) flow cytometry analysis of NCI-H292 cells (bottom histogram) compared to a negative control cell (top histogram).FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

SNRPD2 Antibody (N-term) - Background

SNRPD2 belongs to the small nuclear ribonucleoprotein core protein family. It is required for pre-mRNA splicing and small nuclear ribonucleoprotein biogenesis.

SNRPD2 Antibody (N-term) - References

Stelzl,U., et.al., Cell 122 (6), 957-968 (2005)