

ATP5I Polyclonal Antibody
Catalog # AP68590**Specification**

ATP5I Polyclonal Antibody - Product Information

Application	WB, IHC-P
Primary Accession	P56385
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal

ATP5I Polyclonal Antibody - Additional Information**Gene ID** 521**Other Names**

ATP5I; ATP5K; ATP synthase subunit e; mitochondrial; ATPase subunit e

Dilution

WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/20000. Not yet tested in other applications.

IHC-P~~N/A

Format

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

Storage Conditions

-20°C

ATP5I Polyclonal Antibody - Protein Information**Name** ATP5ME ([HGNC:846](#))**Function**

Subunit e, of the mitochondrial membrane ATP synthase complex (F(1)F(0) ATP synthase or Complex V) that produces ATP from ADP in the presence of a proton gradient across the membrane which is generated by electron transport complexes of the respiratory chain (PubMed:37244256). ATP synthase complex consist of a soluble F(1) head domain - the catalytic core - and a membrane F(1) domain - the membrane proton channel (PubMed:37244256). These two domains are linked by a central stalk rotating inside the F(1) region and a stationary peripheral stalk (PubMed:37244256). During catalysis, ATP synthesis in the catalytic domain of F(1) is coupled via a rotary mechanism of the central stalk subunits to proton translocation (Probable). In vivo, can only synthesize ATP although its ATP hydrolase activity can be activated artificially in vitro (By similarity). Part of the complex F(0) domain (PubMed:37244256).


Cellular Location

Mitochondrion. Mitochondrion inner membrane.

ATP5I Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

ATP5I Polyclonal Antibody - Images**ATP5I Polyclonal Antibody - Background**

Mitochondrial membrane ATP synthase (F₁F₀) ATP synthase or Complex V) produces ATP from ADP in the presence of a proton gradient across the membrane which is generated by electron transport complexes of the respiratory chain. F-type ATPases consist of two structural domains, F₁ - containing the extramembraneous catalytic core, and F₀ - containing the membrane proton channel, linked together by a central stalk and a peripheral stalk. During catalysis, ATP synthesis in the catalytic domain of F₁ is coupled via a rotary mechanism of the central stalk subunits to proton translocation. Part of the complex F₀ domain. Minor subunit located with subunit a in the membrane.