

## **Cathepsin D Polyclonal Antibody**

Catalog # AP68864

## **Specification**

## **Cathepsin D Polyclonal Antibody - Product Information**

Application WB, IHC-P
Primary Accession P07339
Reactivity Human
Host Rabbit
Clonality Polyclonal

## Cathepsin D Polyclonal Antibody - Additional Information

**Gene ID 1509** 

**Other Names** 

CTSD; CPSD; Cathepsin D

#### **Dilution**

WB $\sim\sim$ Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/40000. Not yet tested in other applications.

IHC-P~~N/A

#### **Format**

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

### **Storage Conditions**

-20°C

### **Cathepsin D Polyclonal Antibody - Protein Information**

Name CTSD

**Synonyms** CPSD

### **Function**

Acid protease active in intracellular protein breakdown. Plays a role in APP processing following cleavage and activation by ADAM30 which leads to APP degradation (PubMed:<a href="http://www.uniprot.org/citations/27333034" target="\_blank">27333034</a>). Involved in the pathogenesis of several diseases such as breast cancer and possibly Alzheimer disease.

#### **Cellular Location**

Lysosome. Melanosome. Secreted, extracellular space. Note=Identified by mass spectrometry in melanosome fractions from stage I to stage IV. In aortic samples, detected as an extracellular protein loosely bound to the matrix (PubMed:20551380)

#### **Tissue Location**

Expressed in the aorta extracellular space (at protein level) (PubMed:20551380). Expressed in



Tel: 858.875.1900 Fax: 858.875.1999


liver (at protein level) (PubMed:1426530).

# Cathepsin D Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

## Cathepsin D Polyclonal Antibody - Images



# Cathepsin D Polyclonal Antibody - Background

Acid protease active in intracellular protein breakdown. Plays a role in APP processing following cleavage and activation by ADAM30 which leads to APP degradation (PubMed:27333034). Involved in the pathogenesis of several diseases such as breast cancer and possibly Alzheimer disease.