

**CD71 Polyclonal Antibody**  
**Catalog # AP68958****Specification****CD71 Polyclonal Antibody - Product Information**

Application	WB, IHC-P
Primary Accession	<a href="#">P02786</a>
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal

**CD71 Polyclonal Antibody - Additional Information****Gene ID** 7037**Other Names**

TFR1; Transferrin receptor protein 1; TR; TfR; TfR1; Trfr; T9; p90; CD antigen CD71

**Dilution**

WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/10000. Not yet tested in other applications.

IHC-P~~N/A

**Format**

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

**Storage Conditions**

-20°C

**CD71 Polyclonal Antibody - Protein Information****Name** TFR1**Function**

Cellular uptake of iron occurs via receptor-mediated endocytosis of ligand-occupied transferrin receptor into specialized endosomes (PubMed:<a href="http://www.uniprot.org/citations/26214738" target="\_blank">26214738</a>). Endosomal acidification leads to iron release. The apotransferrin-receptor complex is then recycled to the cell surface with a return to neutral pH and the concomitant loss of affinity of apotransferrin for its receptor. Transferrin receptor is necessary for development of erythrocytes and the nervous system (By similarity). A second ligand, the hereditary hemochromatosis protein HFE, competes for binding with transferrin for an overlapping C- terminal binding site. Positively regulates T and B cell proliferation through iron uptake (PubMed:<a href="http://www.uniprot.org/citations/26642240" target="\_blank">26642240</a>). Acts as a lipid sensor that regulates mitochondrial fusion by regulating activation of the JNK pathway (PubMed:<a href="http://www.uniprot.org/citations/26214738" target="\_blank">26214738</a>). When dietary levels of stearate (C18:0) are low, promotes activation of the JNK pathway, resulting in HUWE1- mediated ubiquitination and subsequent degradation of the mitofusin MFN2 and

inhibition of mitochondrial fusion (PubMed:<a href="http://www.uniprot.org/citations/26214738" target="\_blank">26214738</a>). When dietary levels of stearate (C18:0) are high, TFRC stearylation inhibits activation of the JNK pathway and thus degradation of the mitofusin MFN2 (PubMed:<a href="http://www.uniprot.org/citations/26214738" target="\_blank">26214738</a>). Mediates uptake of NICOL1 into fibroblasts where it may regulate extracellular matrix production (By similarity).

#### Cellular Location


Cell membrane; Single-pass type II membrane protein Melanosome. Note=Identified by mass spectrometry in melanosome fractions from stage I to stage IV

### CD71 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### CD71 Polyclonal Antibody - Images



### CD71 Polyclonal Antibody - Background

Cellular uptake of iron occurs via receptor-mediated endocytosis of ligand-occupied transferrin receptor into specialized endosomes. Endosomal acidification leads to iron release. The apotransferrin-receptor complex is then recycled to the cell surface with a return to neutral pH and the concomitant loss of affinity of apotransferrin for its receptor. Transferrin receptor is necessary for development of erythrocytes and the nervous system (By similarity). A second ligand, the hereditary hemochromatosis protein HFE, competes for binding with transferrin for an overlapping C-terminal binding site. Positively regulates T and B cell proliferation through iron uptake

(PubMed:26642240).