

VCP Antibody (C-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP6920b**Specification**

VCP Antibody (C-term) - Product Information

Application	FC, IHC-P, WB,E
Primary Accession	P55072
Other Accession	P23787 , P46462 , P03974 , Q01853 , Q7ZU99 , Q3ZBT1
Reactivity	Human
Predicted	Bovine, Zebrafish, Mouse, Pig, Rat, Xenopus
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Antigen Region	726-755

VCP Antibody (C-term) - Additional Information**Gene ID** 7415**Other Names**Transitional endoplasmic reticulum ATPase, TER ATPase, 15S Mg(2+)-ATPase p97 subunit,
Valosin-containing protein, VCP, VCP**Target/Specificity**

This VCP antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 726-755 amino acids from the C-terminal region of human VCP.

Dilution

FC~~1:10~50

IHC-P~~1:50~100

WB~~1:1000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

VCP Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

VCP Antibody (C-term) - Protein Information

Name VCP

Function Necessary for the fragmentation of Golgi stacks during mitosis and for their reassembly after mitosis. Involved in the formation of the transitional endoplasmic reticulum (tER). The transfer of membranes from the endoplasmic reticulum to the Golgi apparatus occurs via 50-70 nm transition vesicles which derive from part-rough, part-smooth transitional elements of the endoplasmic reticulum (tER). Vesicle budding from the tER is an ATP-dependent process. The ternary complex containing UFD1, VCP and NPLOC4 binds ubiquitinated proteins and is necessary for the export of misfolded proteins from the ER to the cytoplasm, where they are degraded by the proteasome. The NPLOC4- UFD1-VCP complex regulates spindle disassembly at the end of mitosis and is necessary for the formation of a closed nuclear envelope. Regulates E3 ubiquitin-protein ligase activity of RNF19A. Component of the VCP/p97-AMFR/gp78 complex that participates in the final step of the sterol-mediated ubiquitination and endoplasmic reticulum-associated degradation (ERAD) of HMGCR. Mediates the endoplasmic reticulum- associated degradation of CHRNA3 in cortical neurons as part of the STUB1-VCP-UBXN2A complex (PubMed:[26265139](#)). Involved in endoplasmic reticulum stress-induced pre-emptive quality control, a mechanism that selectively attenuates the translocation of newly synthesized proteins into the endoplasmic reticulum and reroutes them to the cytosol for proteasomal degradation (PubMed:[26565908](#)). Involved in clearance process by mediating G3BP1 extraction from stress granules (PubMed:[29804830](#), PubMed:[34739333](#)). Also involved in DNA damage response: recruited to double-strand breaks (DSBs) sites in a RNF8- and RNF168-dependent manner and promotes the recruitment of TP53BP1 at DNA damage sites (PubMed:[22020440](#), PubMed:[22120668](#)). Recruited to stalled replication forks by SPRTN: may act by mediating extraction of DNA polymerase eta (POLH) to prevent excessive translesion DNA synthesis and limit the incidence of mutations induced by DNA damage (PubMed:[23042605](#), PubMed:[23042607](#)). Together with SPRTN metalloprotease, involved in the repair of covalent DNA-protein cross- links (DPCs) during DNA synthesis (PubMed:[32152270](#)). Involved in interstrand cross-link repair in response to replication stress by mediating unloading of the ubiquitinated CMG helicase complex (By similarity). Mediates extraction of PARP1 trapped to chromatin: recognizes and binds ubiquitinated PARP1 and promotes its removal (PubMed:[35013556](#)). Required for cytoplasmic retrotranslocation of stressed/damaged mitochondrial outer-membrane proteins and their subsequent proteasomal degradation (PubMed:[16186510](#), PubMed:[21118995](#)). Essential for the maturation of ubiquitin-containing autophagosomes and the clearance of ubiquitinated protein by autophagy (PubMed:[20104022](#), PubMed:[27753622](#)). Acts as a negative regulator of type I interferon production by interacting with RIGI: interaction takes place when RIGI is ubiquitinated via 'Lys-63'-linked ubiquitin on its CARD domains, leading to recruit RNF125 and promote ubiquitination and degradation of RIGI (PubMed:[26471729](#)). May play a role in the ubiquitin-dependent sorting of membrane proteins to lysosomes where they undergo degradation (PubMed:[21822278](#)). May more particularly play a role in caveolins sorting in cells (PubMed:[21822278](#), PubMed:[23335559](#)). By controlling the steady-state expression of the IGF1R receptor, indirectly regulates the insulin-like growth factor receptor signaling pathway (PubMed:[26692333](#)).

Cellular Location


Cytoplasm, cytosol. Endoplasmic reticulum. Nucleus. Cytoplasm, Stress granule. Note=Present in the neuronal hyaline inclusion bodies specifically found in motor neurons from amyotrophic lateral sclerosis patients (PubMed:[15456787](#)). Present in the Lewy bodies specifically found in neurons from Parkinson disease patients (PubMed:[15456787](#)). Recruited to the cytoplasmic surface of the endoplasmic reticulum via interaction with AMFR/gp78 (PubMed:[16168377](#)) Following DNA double-strand breaks, recruited to the sites of damage (PubMed:[22120668](#)). Recruited to stalled replication forks via interaction with SPRTN (PubMed:[23042605](#)). Recruited to damaged lysosomes decorated with K48-linked ubiquitin chains (PubMed:[27753622](#)) Colocalizes with TIA1, ZFAND1 and G3BP1 in cytoplasmic stress granules (SGs) in response to arsenite-induced stress treatment (PubMed:[29804830](#)).

VCP Antibody (C-term) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


VCP Antibody (C-term) - Images



Western blot analysis of VCP Antibody (C-term) (Cat. #AP6920b) in Jurkat, 293, MDA-MB231 cell line lysates (35ug/lane). VCP (arrow) was detected using the purified Pab.



Formalin-fixed and paraffin-embedded human brain tissue reacted with VCP Antibody (C-term), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.



Flow cytometric analysis of MDA-231 cells using VCP Antibody (C-term)(bottom histogram) compared to a negative control cell (top histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

VCP Antibody (C-term) - Background

VCP is a member of a family that includes putative ATP-binding proteins involved in vesicle transport and fusion, 26S proteasome function, and assembly of peroxisomes. This protein, as a structural protein, is associated with clathrin, and heat-shock protein Hsc70, to form a complex. It has been implicated in a number of cellular events that are regulated during mitosis, including homotypic membrane fusion, spindle pole body function, and ubiquitin-dependent protein degradation.

VCP Antibody (C-term) - References

Rush,J., et.al., Nat. Biotechnol. 23 (1), 94-101 (2005)

VCP Antibody (C-term) - Citations

- [KUS121, a valosin-containing protein modulator, attenuates ischemic stroke via preventing ATP depletion.](#)
- [Immunoreactivity of valosin-containing protein in sporadic amyotrophic lateral sclerosis and in a case of its novel mutant.](#)