

**cPLA2 Polyclonal Antibody**  
**Catalog # AP69267****Specification**

---

**cPLA2 Polyclonal Antibody - Product Information**

|                   |                        |
|-------------------|------------------------|
| Application       | WB, IHC-P, IF          |
| Primary Accession | <a href="#">P47712</a> |
| Reactivity        | Human, Mouse, Rat      |
| Host              | Rabbit                 |
| Clonality         | Polyclonal             |

**cPLA2 Polyclonal Antibody - Additional Information****Gene ID** 5321**Other Names**

PLA2G4A; CPLA2; PLA2G4; Cytosolic phospholipase A2; cPLA2; Phospholipase A2 group IVA

**Dilution**

WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/10000. Not yet tested in other applications.

IHC-P~~N/A

IF~~1:50~200

**Format**

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

**Storage Conditions**

-20°C

**cPLA2 Polyclonal Antibody - Protein Information****Name** PLA2G4A**Synonyms** CPLA2, PLA2G4**Function**

Has primarily calcium-dependent phospholipase and lysophospholipase activities, with a major role in membrane lipid remodeling and biosynthesis of lipid mediators of the inflammatory response (PubMed:<a href="http://www.uniprot.org/citations/10358058" target="\_blank">10358058</a>, PubMed:<a href="http://www.uniprot.org/citations/14709560" target="\_blank">14709560</a>, PubMed:<a href="http://www.uniprot.org/citations/16617059" target="\_blank">16617059</a>, PubMed:<a href="http://www.uniprot.org/citations/17472963" target="\_blank">17472963</a>, PubMed:<a href="http://www.uniprot.org/citations/18451993" target="\_blank">18451993</a>, PubMed:<a href="http://www.uniprot.org/citations/27642067" target="\_blank">27642067</a>, PubMed:<a href="http://www.uniprot.org/citations/7794891" target="\_blank">7794891</a>, PubMed:<a href="http://www.uniprot.org/citations/8619991" target="\_blank">8619991</a>, PubMed:<a href="http://www.uniprot.org/citations/8702602" target="\_blank">8702602</a>),

PubMed:<a href="http://www.uniprot.org/citations/9425121" target="\_blank">9425121</a>). Plays an important role in embryo implantation and parturition through its ability to trigger prostanoid production (By similarity). Preferentially hydrolyzes the ester bond of the fatty acyl group attached at sn-2 position of phospholipids (phospholipase A2 activity) (PubMed:<a href="http://www.uniprot.org/citations/10358058" target="\_blank">10358058</a>, PubMed:<a href="http://www.uniprot.org/citations/17472963" target="\_blank">17472963</a>, PubMed:<a href="http://www.uniprot.org/citations/18451993" target="\_blank">18451993</a>, PubMed:<a href="http://www.uniprot.org/citations/7794891" target="\_blank">7794891</a>, PubMed:<a href="http://www.uniprot.org/citations/8619991" target="\_blank">8619991</a>, PubMed:<a href="http://www.uniprot.org/citations/9425121" target="\_blank">9425121</a>). Selectively hydrolyzes sn-2 arachidonoyl group from membrane phospholipids, providing the precursor for eicosanoid biosynthesis via the cyclooxygenase pathway (PubMed:<a href="http://www.uniprot.org/citations/10358058" target="\_blank">10358058</a>, PubMed:<a href="http://www.uniprot.org/citations/17472963" target="\_blank">17472963</a>, PubMed:<a href="http://www.uniprot.org/citations/18451993" target="\_blank">18451993</a>, PubMed:<a href="http://www.uniprot.org/citations/7794891" target="\_blank">7794891</a>, PubMed:<a href="http://www.uniprot.org/citations/9425121" target="\_blank">9425121</a>). In an alternative pathway of eicosanoid biosynthesis, hydrolyzes sn-2 fatty acyl chain of eicosanoid lysophospholipids to release free bioactive eicosanoids (PubMed:<a href="http://www.uniprot.org/citations/27642067" target="\_blank">27642067</a>). Hydrolyzes the ester bond of the fatty acyl group attached at sn-1 position of phospholipids (phospholipase A1 activity) only if an ether linkage rather than an ester linkage is present at the sn-2 position. This hydrolysis is not stereospecific (PubMed:<a href="http://www.uniprot.org/citations/7794891" target="\_blank">7794891</a>). Has calcium-independent phospholipase A2 and lysophospholipase activities in the presence of phosphoinositides (PubMed:<a href="http://www.uniprot.org/citations/12672805" target="\_blank">12672805</a>). Has O-acyltransferase activity. Catalyzes the transfer of fatty acyl chains from phospholipids to a primary hydroxyl group of glycerol (sn-1 or sn-3), potentially contributing to monoacylglycerol synthesis (PubMed:<a href="http://www.uniprot.org/citations/7794891" target="\_blank">7794891</a>).

#### **Cellular Location**

Cytoplasm. Golgi apparatus membrane. Nucleus envelope Note=Translocates to intracellular membranes in a calcium-dependent way.

#### **Tissue Location**


Expressed in various cells and tissues such as macrophages, neutrophils, fibroblasts and lung endothelium. Expressed in platelets (at protein level) (PubMed:25102815)


### **cPLA2 Polyclonal Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### **cPLA2 Polyclonal Antibody - Images**





#### **cPLA2 Polyclonal Antibody - Background**

Selectively hydrolyzes arachidonyl phospholipids in the sn-2 position releasing arachidonic acid. Together with its lysophospholipid activity, it is implicated in the initiation of the inflammatory response.