

Creatine Kinase M Polyclonal Antibody
Catalog # AP69279**Specification**

Creatine Kinase M Polyclonal Antibody - Product Information

Application	WB, IHC-P, IF
Primary Accession	P06732
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal

Creatine Kinase M Polyclonal Antibody - Additional Information**Gene ID** 1158**Other Names**

CKM; CKMM; Creatine kinase M-type; Creatine kinase M chain; M-CK

Dilution

WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/40000. Not yet tested in other applications.

IHC-P~~N/A

IF~~1:50~200

Format

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

Storage Conditions

-20°C

Creatine Kinase M Polyclonal Antibody - Protein Information**Name** CKM**Synonyms** CKMM**Function**

Reversibly catalyzes the transfer of phosphate between ATP and various phosphogens (e.g. creatine phosphate). Creatine kinase isoenzymes play a central role in energy transduction in tissues with large, fluctuating energy demands, such as skeletal muscle, heart, brain and spermatozoa.

Cellular Location



Cytoplasm.


Creatine Kinase M Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Creatine Kinase M Polyclonal Antibody - Images





Creatine Kinase M Polyclonal Antibody - Background

Reversibly catalyzes the transfer of phosphate between ATP and various phosphogens (e.g. creatine phosphate). Creatine kinase isoenzymes play a central role in energy transduction in tissues with large, fluctuating energy demands, such as skeletal muscle, heart, brain and spermatozoa.