

CTDSP1 Polyclonal Antibody
Catalog # AP69333**Specification**

CTDSP1 Polyclonal Antibody - Product Information

Application	WB, IHC-P
Primary Accession	Q9GZU7
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal

CTDSP1 Polyclonal Antibody - Additional Information**Gene ID** 58190**Other Names**

CTDSP1; NIF3; NLIIF; SCP1; Carboxy-terminal domain RNA polymerase II polypeptide A small phosphatase 1; Nuclear LIM interactor-interacting factor 3; NLI-IF; NLI-interacting factor 3; Small C-terminal domain phosphatase 1; SCP1; Small CTD ph

Dilution

WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/40000. Not yet tested in other applications.
IHC-P~~N/A

Format

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

Storage Conditions

-20°C

CTDSP1 Polyclonal Antibody - Protein Information**Name** CTDSP1**Synonyms** NIF3, NLIIF, SCP1**Function**

Preferentially catalyzes the dephosphorylation of 'Ser-5' within the tandem 7 residue repeats in the C-terminal domain (CTD) of the largest RNA polymerase II subunit POLR2A. Negatively regulates RNA polymerase II transcription, possibly by controlling the transition from initiation/capping to processive transcript elongation. Recruited by REST to neuronal genes that contain RE-1 elements, leading to neuronal gene silencing in non-neuronal cells.

Cellular Location

Nucleus. Note=Colocalizes with RNA polymerase II

Tissue Location



Expression is restricted to non-neuronal tissues. Highest expression in skeletal muscle, spleen, lung and placenta

CTDSP1 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

CTDSP1 Polyclonal Antibody - Images



CTDSP1 Polyclonal Antibody - Background

Preferentially catalyzes the dephosphorylation of 'Ser- 5' within the tandem 7 residue repeats in the C-terminal domain (CTD) of the largest RNA polymerase II subunit POLR2A. Negatively regulates RNA polymerase II transcription, possibly by controlling the transition from initiation/capping to processive transcript elongation. Recruited by REST to neuronal genes that contain RE-1 elements, leading to neuronal gene silencing in non-neuronal cells.