

Eg5 Polyclonal Antibody
Catalog # AP69663**Specification**

Eg5 Polyclonal Antibody - Product Information

Application	WB, IHC-P, IF
Primary Accession	P52732
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal

Eg5 Polyclonal Antibody - Additional Information**Gene ID** 3832**Other Names**

KIF11; EG5; KNSL1; TRIP5; Kinesin-like protein KIF11; Kinesin-like protein 1; Kinesin-like spindle protein HKSP; Kinesin-related motor protein Eg5; Thyroid receptor-interacting protein 5; TR-interacting protein 5; TRIP-5

Dilution

WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/20000. Not yet tested in other applications.
IHC-P~~N/A
IF~~1:50~200

Format

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

Storage Conditions

-20°C

Eg5 Polyclonal Antibody - Protein Information**Name** KIF11**Synonyms** EG5, KNSL1, TRIP5**Function**

Motor protein required for establishing a bipolar spindle and thus contributing to chromosome congression during mitosis (PubMed:19001501, PubMed:37728657). Required in non-mitotic cells for transport of secretory proteins from the Golgi complex to the cell surface (PubMed:23857769).

Cellular Location


Cytoplasm. Cytoplasm, cytoskeleton, spindle pole


Eg5 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Eg5 Polyclonal Antibody - Images





Eg5 Polyclonal Antibody - Background

Motor protein required for establishing a bipolar spindle during mitosis (PubMed:19001501). Required in non-mitotic cells for transport of secretory proteins from the Golgi complex to the cell surface (PubMed:23857769).