

MMP-19 Polyclonal Antibody
Catalog # AP70980**Specification**

MMP-19 Polyclonal Antibody - Product Information

Application	WB, IHC-P
Primary Accession	Q99542
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal

MMP-19 Polyclonal Antibody - Additional Information**Gene ID** 4327**Other Names**MMP19; MMP18; RASI; Matrix metalloproteinase-19; MMP-19; Matrix metalloproteinase RASI;
Matrix metalloproteinase-18; MMP-18**Dilution**WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/10000. Not
yet tested in other applications.

IHC-P~~N/A

Format

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

Storage Conditions

-20°C

MMP-19 Polyclonal Antibody - Protein Information**Name** MMP19**Synonyms** MMP18, RASI**Function**Endopeptidase that degrades various components of the extracellular matrix, such as aggrecan
and cartilage oligomeric matrix protein (comp), during development, haemostasis and pathological
conditions (arthritic disease). May also play a role in neovascularization or angiogenesis.
Hydrolyzes collagen type IV, laminin, nidogen, nascin-C isoform, fibronectin, and type I gelatin.**Cellular Location**

Secreted, extracellular space, extracellular matrix

Tissue LocationExpressed in mammary gland, placenta, lung, pancreas, ovary, small intestine, spleen, thymus,
prostate, testis colon, heart and blood vessel walls. Not detected in brain and peripheral blood


leukocytes. Also expressed in the synovial fluid of normal and rheumatoid patients (PubMed:8920941)

MMP-19 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

MMP-19 Polyclonal Antibody - Images



MMP-19 Polyclonal Antibody - Background

Endopeptidase that degrades various components of the extracellular matrix, such as aggrecan and cartilage oligomeric matrix protein (comp), during development, haemostasis and pathological conditions (arthritic disease). May also play a role in neovascularization or angiogenesis. Hydrolyzes collagen type IV, laminin, nidogen, nascin-C isoform, fibronectin, and type I gelatin.