

P2RY8 Polyclonal Antibody
Catalog # AP71687**Specification****P2RY8 Polyclonal Antibody - Product Information**

Application	WB, IF
Primary Accession	Q86VZ1
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal

P2RY8 Polyclonal Antibody - Additional Information**Gene ID** 286530**Other Names**

P2RY8; P2Y purinoceptor 8; P2Y8

Dilution

WB~~Western Blot: 1/500 - 1/2000. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/20000. Not yet tested in other applications.

IF~~1:50~200

Format

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

Storage Conditions

-20°C

P2RY8 Polyclonal Antibody - Protein Information**Name** P2RY8 {ECO:0000303|PubMed:30842656, ECO:0000312|HGNC:HGNC:15524}**Function**

G protein-coupled receptor for S-geranylgeranyl-glutathione (GGG), an endogenous metabolite present in lymphoid tissues. Couples the binding of GGG to the activation of GNA13 and downstream repression of AKT activation in lymphocytes defining their positioning and growth within lymphoid organs (PubMed:25274307, PubMed:30842656, PubMed:34088745). In lymphoid follicles, confines B cells and follicular helper T cells in germinal centers (GCs) in response to GGG local gradients established by GGT5 (via GGG catabolism) and ABCC1 (via extracellular transport) with lower concentrations of GGG found in the follicular dendritic cell network region around which germinal centers are formed (PubMed:25274307, PubMed:30842656, PubMed:34088745). In the bone marrow, also in response to GGG gradients established by GGT5 and ABCC1, it restricts

chemotactic transmigration of B cells, T cells and NK cells from blood vessels to the bone marrow parenchyma (PubMed:30842656, PubMed:34088745). Contributes to GNA13-dependent pathway that suppresses GC B cell growth (PubMed:25274307).

Cellular Location

Cell membrane; Multi-pass membrane protein

Tissue Location


Barely detectable in normal blood leukocytes. Weaker expression was seen in heart, kidney and lung. Not detected in brain (PubMed:11004484, PubMed:15466006). Expressed in B cells and follicular helper T cells in germinal centers (at protein level) (PubMed:30842656).


P2RY8 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

P2RY8 Polyclonal Antibody - Images





P2RY8 Polyclonal Antibody - Background

Probable receptor for purines coupled to G-proteins.