

**Keap1 Polyclonal Antibody**  
**Catalog # AP73401****Specification****Keap1 Polyclonal Antibody - Product Information**

|                   |                        |
|-------------------|------------------------|
| Application       | WB, IHC-P              |
| Primary Accession | <a href="#">Q14145</a> |
| Reactivity        | Human, Mouse, Rat      |
| Host              | Rabbit                 |
| Clonality         | Polyclonal             |

**Keap1 Polyclonal Antibody - Additional Information****Gene ID** 9817**Other Names**

KEAP1; INRF2; KIAA0132; KLHL19; Kelch-like ECH-associated protein 1; Cytosolic inhibitor of Nrf2; INrf2; Kelch-like protein 19

**Dilution**

WB~~Western Blot: 1/500 - 1/2000. IHC-p: 1:100-300 ELISA: 1/20000. Not yet tested in other applications.

IHC-P~~N/A

**Format**

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

**Storage Conditions**

-20°C

**Keap1 Polyclonal Antibody - Protein Information****Name** KEAP1 {ECO:0000303|PubMed:14585973, ECO:0000312|HGNC:HGNC:23177}**Function**

Substrate-specific adapter of a BCR (BTB-CUL3-RBX1) E3 ubiquitin ligase complex that regulates the response to oxidative stress by targeting NFE2L2/NRF2 for ubiquitination (PubMed:<a href="http://www.uniprot.org/citations/14585973" target="\_blank">14585973</a>, PubMed:<a href="http://www.uniprot.org/citations/15379550" target="\_blank">15379550</a>, PubMed:<a href="http://www.uniprot.org/citations/15572695" target="\_blank">15572695</a>, PubMed:<a href="http://www.uniprot.org/citations/15601839" target="\_blank">15601839</a>, PubMed:<a href="http://www.uniprot.org/citations/15983046" target="\_blank">15983046</a>, PubMed:<a href="http://www.uniprot.org/citations/37339955" target="\_blank">37339955</a>). KEAP1 acts as a key sensor of oxidative and electrophilic stress: in normal conditions, the BCR(KEAP1) complex mediates ubiquitination and degradation of NFE2L2/NRF2, a transcription factor regulating expression of many cytoprotective genes (PubMed:<a href="http://www.uniprot.org/citations/15601839" target="\_blank">15601839</a>, PubMed:<a href="http://www.uniprot.org/citations/16006525" target="\_blank">16006525</a>). In response

to oxidative stress, different electrophile metabolites trigger non-enzymatic covalent modifications of highly reactive cysteine residues in KEAP1, leading to inactivate the ubiquitin ligase activity of the BCR(KEAP1) complex, promoting NFE2L2/NRF2 nuclear accumulation and expression of phase II detoxifying enzymes (PubMed:<a href="http://www.uniprot.org/citations/16006525" target="\_blank">16006525</a>, PubMed:<a href="http://www.uniprot.org/citations/17127771" target="\_blank">17127771</a>, PubMed:<a href="http://www.uniprot.org/citations/18251510" target="\_blank">18251510</a>, PubMed:<a href="http://www.uniprot.org/citations/19489739" target="\_blank">19489739</a>, PubMed:<a href="http://www.uniprot.org/citations/29590092" target="\_blank">29590092</a>). In response to selective autophagy, KEAP1 is sequestered in inclusion bodies following its interaction with SQSTM1/p62, leading to inactivation of the BCR(KEAP1) complex and activation of NFE2L2/NRF2 (PubMed:<a href="http://www.uniprot.org/citations/20452972" target="\_blank">20452972</a>). The BCR(KEAP1) complex also mediates ubiquitination of SQSTM1/p62, increasing SQSTM1/p62 sequestering activity and degradation (PubMed:<a href="http://www.uniprot.org/citations/28380357" target="\_blank">28380357</a>). The BCR(KEAP1) complex also targets BPTF and PGAM5 for ubiquitination and degradation by the proteasome (PubMed:<a href="http://www.uniprot.org/citations/15379550" target="\_blank">15379550</a>, PubMed:<a href="http://www.uniprot.org/citations/17046835" target="\_blank">17046835</a>).

#### **Cellular Location**

Cytoplasm. Nucleus. Note=Mainly cytoplasmic (PubMed:15601839). In response to selective autophagy, relocates to inclusion bodies following interaction with SQSTM1/p62 (PubMed:20452972).

#### **Tissue Location**


Broadly expressed, with highest levels in skeletal muscle.




### **Keap1 Polyclonal Antibody - Protocols**




Provided below are standard protocols that you may find useful for product applications.





- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### **Keap1 Polyclonal Antibody - Images**









**Keap1 Polyclonal Antibody - Background**

Acts as a substrate adapter protein for the E3 ubiquitin ligase complex formed by CUL3 and RBX1 and targets NFE2L2/NRF2 for ubiquitination and degradation by the proteasome, thus resulting in the suppression of its transcriptional activity and the repression of antioxidant response element-mediated detoxifying enzyme gene expression. Retains NFE2L2/NRF2 and may also retain BPTF in the cytosol. Targets PGAM5 for ubiquitination and degradation by the proteasome.