

## Specification

### CD73 Polyclonal Antibody - Additional Information

**Gene ID 4907**

## Other Names

NT5E: NT5: NTE: 5'-nucleotidase: 5'-NT: Ecto-5'-nucleotidase: CD73

## Dilution

WB~~Western Blot: 1/500 - 1/2000. IHC-p: 1:100-300 ELISA: 1/20000. Not yet tested in other applications.

IHC-P ~ ~ N/A

## Format

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

## Storage Conditions

-20°C

## CD73 Polyclonal Antibody - Protein Information

Name NT5E

**Synonyms** NT5, NTE

## Function

Catalyzes the hydrolysis of nucleotide monophosphates, releasing inorganic phosphate and the corresponding nucleoside, with AMP being the preferred substrate (PubMed:<a href="http://www.uniprot.org/citations/21933152" target="\_blank">21933152</a>, PubMed:<a href="http://www.uniprot.org/citations/22997138" target="\_blank">22997138</a>, PubMed:<a href="http://www.uniprot.org/citations/23142347" target="\_blank">23142347</a>, PubMed:<a href="http://www.uniprot.org/citations/24887587" target="\_blank">24887587</a>, PubMed:<a href="http://www.uniprot.org/citations/34403084" target="\_blank">34403084</a>). Shows a preference for ribonucleotide monophosphates over their equivalent deoxyribose forms (PubMed:<a href="http://www.uniprot.org/citations/34403084" target="\_blank">34403084</a>). Other substrates include IMP, UMP, GMP, CMP, dAMP, dCMP, dTMP, NAD and NMN (PubMed:<a href="http://www.uniprot.org/citations/21933152" target="\_blank">21933152</a>, PubMed:<a href="http://www.uniprot.org/citations/22997138" target="\_blank">22997138</a>, PubMed:<a href="http://www.uniprot.org/citations/23142347" target="\_blank">23142347</a>, PubMed:<a href="http://www.uniprot.org/citations/24887587" target="\_blank">24887587</a>, PubMed:<a href="http://www.uniprot.org/citations/34403084" target="\_blank">34403084</a>).

href="http://www.uniprot.org/citations/23142347" target="\_blank">23142347</a>, PubMed:<a href="http://www.uniprot.org/citations/24887587" target="\_blank">24887587</a>, PubMed:<a href="http://www.uniprot.org/citations/34403084" target="\_blank">34403084</a>).

#### Cellular Location



Cell membrane; Lipid-anchor, GPI-anchor

#### CD73 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### CD73 Polyclonal Antibody - Images



**CD73 Polyclonal Antibody - Background**

Hydrolyzes extracellular nucleotides into membrane permeable nucleosides. Exhibits AMP-, NAD-, and NMN-nucleosidase activities.