

ENA-78 Polyclonal Antibody
Catalog # AP74144**Specification**

ENA-78 Polyclonal Antibody - Product Information

Application	IHC-P
Primary Accession	P42830
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal

ENA-78 Polyclonal Antibody - Additional Information**Gene ID** 6374**Other Names**

C-X-C motif chemokine 5 (ENA-78(1-78)) (Epithelial-derived neutrophil-activating protein 78) (Neutrophil-activating peptide ENA-78) (Small-inducible cytokine B5) [Cleaved into: ENA-78(8-78); ENA-78(9-78)]

Dilution

IHC-P~~N/A

Format

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

Storage Conditions

-20°C

ENA-78 Polyclonal Antibody - Protein Information**Name** CXCL5**Synonyms** ENA78, SCYB5**Function**

Involved in neutrophil activation. In vitro, ENA-78(8-78) and ENA-78(9-78) show a threefold higher chemotactic activity for neutrophil granulocytes.

Cellular Location




Secreted.


ENA-78 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

ENA-78 Polyclonal Antibody - Images





ENA-78 Polyclonal Antibody - Background

Involved in neutrophil activation. In vitro, ENA-78(8-78) and ENA-78(9-78) show a threefold higher chemotactic activity for neutrophil granulocytes.