

**VHL Polyclonal Antibody**  
**Catalog # AP74171****Specification**

---

**VHL Polyclonal Antibody - Product Information**

|                   |                        |
|-------------------|------------------------|
| Application       | IHC-P                  |
| Primary Accession | <a href="#">P40337</a> |
| Reactivity        | Human                  |
| Host              | Rabbit                 |
| Clonality         | Polyclonal             |

**VHL Polyclonal Antibody - Additional Information****Gene ID** 7428**Other Names**

Von Hippel-Lindau disease tumor suppressor (Protein G7) (pVHL)

**Dilution**

IHC-P~~N/A

**Format**

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

**Storage Conditions**

-20°C

**VHL Polyclonal Antibody - Protein Information****Name** VHL**Function**

Involved in the ubiquitination and subsequent proteasomal degradation via the von Hippel-Lindau ubiquitination complex (PubMed:<a href="http://www.uniprot.org/citations/10944113" target="\_blank">10944113</a>, PubMed:<a href="http://www.uniprot.org/citations/17981124" target="\_blank">17981124</a>, PubMed:<a href="http://www.uniprot.org/citations/19584355" target="\_blank">19584355</a>). Seems to act as a target recruitment subunit in the E3 ubiquitin ligase complex and recruits hydroxylated hypoxia-inducible factor (HIF) under normoxic conditions (PubMed:<a href="http://www.uniprot.org/citations/10944113" target="\_blank">10944113</a>, PubMed:<a href="http://www.uniprot.org/citations/17981124" target="\_blank">17981124</a>). Involved in transcriptional repression through interaction with HIF1A, HIF1AN and histone deacetylases (PubMed:<a href="http://www.uniprot.org/citations/10944113" target="\_blank">10944113</a>, PubMed:<a href="http://www.uniprot.org/citations/17981124" target="\_blank">17981124</a>). Ubiquitinates, in an oxygen-responsive manner, ADRB2 (PubMed:<a href="http://www.uniprot.org/citations/19584355" target="\_blank">19584355</a>). Acts as a negative regulator of mTORC1 by promoting ubiquitination and degradation of RPTOR (PubMed:<a href="http://www.uniprot.org/citations/34290272" target="\_blank">34290272</a>).

**Cellular Location**

[Isoform 1]: Cytoplasm. Cell membrane; Peripheral membrane protein. Endoplasmic reticulum. Nucleus. Note=Found predominantly in the cytoplasm and with less amounts nuclear or membrane-associated (PubMed:9751722) Colocalizes with ADRB2 at the cell membrane (PubMed:19584355)



**Tissue Location**

Expressed in the adult and fetal brain and kidney.

**VHL Polyclonal Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**VHL Polyclonal Antibody - Images****VHL Polyclonal Antibody - Background**

Involved in the ubiquitination and subsequent proteasomal degradation via the von Hippel-Lindau ubiquitination complex. Seems to act as a target recruitment subunit in the E3 ubiquitin ligase complex and recruits hydroxylated hypoxia- inducible factor (HIF) under normoxic conditions. Involved in transcriptional repression through interaction with HIF1A, HIF1AN and histone deacetylases. Ubiquitinates, in an oxygen-responsive manner, ADRB2.