

MIG Polyclonal Antibody
Catalog # AP74191**Specification**

MIG Polyclonal Antibody - Product Information

Application	IHC-P
Primary Accession	Q07325
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal

MIG Polyclonal Antibody - Additional Information**Gene ID** 4283**Other Names**

C-X-C motif chemokine 9 (Gamma-interferon-induced monokine) (Monokine induced by interferon-gamma) (HuMIG) (MIG) (Small-inducible cytokine B9)

Dilution

IHC-P~~N/A

Format

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

Storage Conditions

-20°C

MIG Polyclonal Antibody - Protein Information**Name** CXCL9**Synonyms** CMK, MIG, SCYB9**Function**

Cytokine that affects the growth, movement, or activation state of cells that participate in immune and inflammatory response. Chemotactic for activated T-cells. Binds to CXCR3.

Cellular Location

Secreted.




MIG Polyclonal Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)

- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

MIG Polyclonal Antibody - Images





MIG Polyclonal Antibody - Background

Cytokine that affects the growth, movement, or activation state of cells that participate in immune and inflammatory response. Chemotactic for activated T-cells. Binds to CXCR3.