

Dlx-2 Polyclonal Antibody
Catalog # AP74264**Specification**

Dlx-2 Polyclonal Antibody - Product Information

Application	WB
Primary Accession	Q07687
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal

Dlx-2 Polyclonal Antibody - Additional Information**Gene ID** 1746**Other Names**

Homeobox protein DLX-2

Dilution

WB~~WB 1:500-2000, ELISA 1:10000-20000

Format

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

Storage Conditions

-20°C

Dlx-2 Polyclonal Antibody - Protein Information**Name** DLX2**Function**

Acts as a transcriptional activator (By similarity). Activates transcription of CGA/alpha-GSU, via binding to the downstream activin regulatory element (DARE) in the gene promoter (By similarity). Plays a role in terminal differentiation of interneurons, such as amacrine and bipolar cells in the developing retina. Likely to play a regulatory role in the development of the ventral forebrain (By similarity). May play a role in craniofacial patterning and morphogenesis (By similarity).

Cellular Location

Nucleus.



Dlx-2 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)

- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Dlx-2 Polyclonal Antibody - Images



Dlx-2 Polyclonal Antibody - Background

Acts as a transcriptional activator. Plays a role in terminal differentiation of interneurons, such as amacrine and bipolar cells in the developing retina. Likely to play a regulatory role in the development of the ventral forebrain. May play a role in craniofacial patterning and morphogenesis.