

DCXR Antibody (Center)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP7443c**Specification**

DCXR Antibody (Center) - Product Information

Application	FC, IHC-P, WB,E
Primary Accession	Q7Z4W1
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	25913
Antigen Region	79-106

DCXR Antibody (Center) - Additional Information**Gene ID** 51181**Other Names**

L-xylulose reductase, XR, Carbonyl reductase II, Dicarboxyl/L-xylulose reductase, Kidney dicarboxyl reductase, kiDCR, Sperm surface protein P34H, DCXR

Target/Specificity

This DCXR antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 79-106 amino acids from the Central region of human DCXR.

Dilution

FC~~1:10~50

IHC-P~~1:50~100

WB~~1:1000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

DCXR Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

DCXR Antibody (Center) - Protein Information**Name** DCXR

Synonyms SDR20C1

Function Catalyzes the NADPH-dependent reduction of several pentoses, tetroses, trioses, alpha-dicarbonyl compounds and L-xylulose. Participates in the uronate cycle of glucose metabolism. May play a role in the water absorption and cellular osmoregulation in the proximal renal tubules by producing xylitol, an osmolyte, thereby preventing osmolytic stress from occurring in the renal tubules.

Cellular Location

Membrane; Peripheral membrane protein. Note=Probably recruited to membranes via an interaction with phosphatidylinositol.


Tissue Location

Highly expressed in kidney, liver and epididymis. In the epididymis, it is mainly expressed in the proximal and distal sections of the corpus region. Weakly or not expressed in brain, lung, heart, spleen and testis.


DCXR Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

DCXR Antibody (Center) - Images

Western blot analysis of anti-DCXR Antibody (Center)(Cat.#AP7443c) in mouse lung tissue lysates (35ug/lane). DCXR (arrow) was detected using the purified Pab.



Formalin-fixed and paraffin-embedded human kidney carcinoma with DCXR Antibody (Center), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.



Flow cytometric analysis of NCI-H292 cells using DCXR Antibody (Center)(bottom histogram) compared to a negative control cell (top histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

DCXR Antibody (Center) - Background

DCXR is an enzyme that has both diacetyl reductase and L-xylulose reductase activities.

DCXR Antibody (Center) - References

- Nakagawa J., Ishikura S.J. Biol. Chem. 277:17883-17891(2002)
- Xia X.Y., Xu X.F., Gao Y.Zhonghua Nan Ke Xue 9:24-27(2003)
- El-Kabbani O., Chung R.P.-T.Acta Crystallogr. D 58:1379-1380(2002)