

BTF3 Rabbit mAb
Catalog # AP75170**Specification**

BTF3 Rabbit mAb - Product Information

Application	WB, IHC-P, IP
Primary Accession	P20290
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Monoclonal Antibody
Calculated MW	22168

BTF3 Rabbit mAb - Additional Information**Gene ID** 689**Other Names**
BTF3**Dilution**
WB~~1/500-1/1000
IHC-P~~N/A
IP~~N/A**Format**
Liquid**BTF3 Rabbit mAb - Protein Information****Name** BTF3**Synonyms** NACB**Function**

When associated with NACA, prevents inappropriate targeting of non-secretory polypeptides to the endoplasmic reticulum (ER). Binds to nascent polypeptide chains as they emerge from the ribosome and blocks their interaction with the signal recognition particle (SRP), which normally targets nascent secretory peptides to the ER. BTF3 is also a general transcription factor that can form a stable complex with RNA polymerase II. Required for the initiation of transcription.

Cellular Location

Cytoplasm. Nucleus. Note=The heterodimer with NACA is cytoplasmic

BTF3 Rabbit mAb - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

BTF3 Rabbit mAb - Images

