

CDC42 Rabbit mAb
Catalog # AP75239**Specification**

CDC42 Rabbit mAb - Product Information

Application	WB
Primary Accession	P60953
Reactivity	Human, Rat
Host	Rabbit
Clonality	Monoclonal Antibody
Calculated MW	21259

CDC42 Rabbit mAb - Additional Information**Gene ID** 998**Other Names**
CDC42**Dilution**
WB~~1/500-1/1000**Format**
Liquid**CDC42 Rabbit mAb - Protein Information****Name** CDC42 ([HGNC:1736](#))**Function**

Plasma membrane-associated small GTPase which cycles between an active GTP-bound and an inactive GDP-bound state. In active state binds to a variety of effector proteins to regulate cellular responses. Involved in epithelial cell polarization processes. Regulates the bipolar attachment of spindle microtubules to kinetochores before chromosome congression in metaphase (PubMed:[15642749](http://www.uniprot.org/citations/15642749)). Regulates cell migration (PubMed:[17038317](http://www.uniprot.org/citations/17038317), PubMed:[22843693](http://www.uniprot.org/citations/22843693)). In neurons, plays a role in the extension and maintenance of the formation of filopodia, thin and actin-rich surface projections (PubMed:[14978216](http://www.uniprot.org/citations/14978216)). Required for DOCK10-mediated spine formation in Purkinje cells and hippocampal neurons. In podocytes, facilitates filopodia and podosomes formation upon DOCK11-activation (PubMed:[33523862](http://www.uniprot.org/citations/33523862)). Upon activation by CaMKII, modulates dendritic spine structural plasticity by relaying CaMKII transient activation to synapse-specific, long-term signaling (By similarity). Also plays a role in phagocytosis through organization of the F-actin cytoskeleton associated with forming phagocytic cups (PubMed:[26465210](http://www.uniprot.org/citations/26465210)). Upon activation by PLEKHG4B, involved in actin cytoskeletal remodeling during epithelial cell-cell

junction formation (PubMed:33310911).

Cellular Location

Cell membrane; Lipid-anchor; Cytoplasmic side. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Cytoplasm, cytoskeleton, spindle. Midbody Cell projection, dendrite {ECO:0000250|UniProtKB:P60766} Note=Localizes to spindle during prometaphase cells. Moves to the central spindle as cells progressed through anaphase to telophase (PubMed:15642749). Localizes at the end of cytokinesis in the intercellular bridge formed between two daughter cells (PubMed:15642749). Its localization is regulated by the activities of guanine nucleotide exchange factor ECT2 and GTPase activating protein RACGAP1 (PubMed:15642749). Colocalizes with NEK6 in the centrosome (PubMed:20873783). In its active GTP-bound form localizes to the leading edge membrane of migrating dendritic cells (By similarity) {ECO:0000250|UniProtKB:P60766, ECO:0000269|PubMed:15642749, ECO:0000269|PubMed:20873783}

CDC42 Rabbit mAb - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

CDC42 Rabbit mAb - Images

