

#### KAT2A Rabbit mAb

**Catalog # AP75480** 

### **Specification**

### **KAT2A Rabbit mAb - Product Information**

Application
Primary Accession
Reactivity
Host
Clonality

WB, IP
092830
Human
Rabbit
Monoclon

**Monoclonal Antibody** 

93926

#### KAT2A Rabbit mAb - Additional Information

**Gene ID 2648** 

Calculated MW

Other Names KAT2A

**Dilution**WB~~1/500-1/1000
IP~~N/A

Format Liquid

# KAT2A Rabbit mAb - Protein Information

Name KAT2A {ECO:0000303|PubMed:27796307, ECO:0000312|HGNC:HGNC:4201}

# **Function**

Protein lysine acyltransferase that can act as a acetyltransferase, glutaryltransferase, succinyltransferase or malonyltransferase, depending on the context (PubMed:<a href="http://www.uniprot.org/citations/29211711" target=" blank">29211711</a>, PubMed:<a href="http://www.uniprot.org/citations/35995428" target=" blank">35995428</a>). Acts as a histone lysine succinyltransferase: catalyzes succinylation of histone H3 on 'Lys-79' (H3K79succ), with a maximum frequency around the transcription start sites of genes (PubMed: <a href="http://www.uniprot.org/citations/29211711" target="\_blank">29211711</a>). Succinylation of histones gives a specific tag for epigenetic transcription activation (PubMed:<a href="http://www.uniprot.org/citations/29211711" target=" blank">29211711</a>). Association with the 2-oxoglutarate dehydrogenase complex, which provides succinyl-CoA, is required for histone succinylation (PubMed:<a href="http://www.uniprot.org/citations/29211711" target=" blank">29211711</a>). In different complexes, functions either as an acetyltransferase (HAT) or as a succinyltransferase: in the SAGA and ATAC complexes, acts as a histone acetyltransferase (PubMed:<a href="http://www.uniprot.org/citations/17301242" target="\_blank">17301242</a>, PubMed:<a href="http://www.uniprot.org/citations/19103755" target="blank">19103755</a>, PubMed:<a href="http://www.uniprot.org/citations/29211711" target="blank">29211711</a>). Has significant histone acetyltransferase activity with core



histones, but not with nucleosome core particles (PubMed: <a href="http://www.uniprot.org/citations/17301242" target=" blank">17301242</a>, PubMed:<a href="http://www.uniprot.org/citations/19103755" target="blank">19103755</a>, PubMed:<a href="http://www.uniprot.org/citations/21131905" target="blank">21131905</a>). Has a a strong preference for acetylation of H3 at 'Lys-9' (H3K9ac) (PubMed:<a href="http://www.uniprot.org/citations/21131905" target=" blank">21131905</a>). Acetylation of histones gives a specific tag for epigenetic transcription activation (PubMed:<a href="http://www.uniprot.org/citations/17301242" target=" blank">17301242</a>, PubMed:<a href="http://www.uniprot.org/citations/19103755" target="\_blank">19103755</a>, PubMed:<a href="http://www.uniprot.org/citations/29211711" target="\_blank">29211711</a>). Recruited by the XPC complex at promoters, where it specifically mediates acetylation of histone variant H2A.Z.1/H2A.Z, thereby promoting expression of target genes (PubMed: <a href="http://www.uniprot.org/citations/29973595" target=" blank">29973595</a>, PubMed:<a href="http://www.uniprot.org/citations/31527837" target=" blank">31527837</a>). Involved in long-term memory consolidation and synaptic plasticity: acts by promoting expression of a hippocampal gene expression network linked to neuroactive receptor signaling (By similarity). Acts as a positive regulator of T-cell activation: upon TCR stimulation, recruited to the IL2 promoter following interaction with NFATC2 and catalyzes acetylation of histone H3 at 'Lys-9' (H3K9ac), leading to promote IL2 expression (By similarity). Required for growth and differentiation of craniofacial cartilage and bone by regulating acetylation of histone H3 at 'Lys-9' (H3K9ac) (By similarity). Regulates embryonic stem cell (ESC) pluripotency and differentiation (By similarity). Also acetylates non- histone proteins, such as CEBPB, MRE11, PPARGC1A, PLK4 and TBX5 (PubMed: <a href="http://www.uniprot.org/citations/16753578" target="blank">16753578</a>, PubMed:<a href="http://www.uniprot.org/citations/17301242" target="\_blank">17301242</a>, PubMed:<a href="http://www.uniprot.org/citations/27796307" target="\_blank">27796307</a>, PubMed:<a href="http://www.uniprot.org/citations/29174768" target="\_blank">29174768</a>, PubMed:<a href="http://www.uniprot.org/citations/38128537" target="blank">38128537</a>). Involved in heart and limb development by mediating acetylation of TBX5, acetylation regulating nucleocytoplasmic shuttling of TBX5 (PubMed:<a href="http://www.uniprot.org/citations/29174768" target=" blank">29174768</a>). Acts as a negative regulator of centrosome amplification by mediating acetylation of PLK4 (PubMed: <a href="http://www.uniprot.org/citations/27796307" target="blank">27796307</a>). Acts as a negative regulator of gluconeogenesis by mediating acetylation and subsequent inactivation of PPARGC1A (PubMed:<a href="http://www.uniprot.org/citations/16753578" target=" blank">16753578</a>, PubMed:<a href="http://www.uniprot.org/citations/23142079" target="blank">23142079</a>). Also acts as a histone glutaryltransferase: catalyzes glutarylation of histone H4 on 'Lys-91' (H4K91glu), a mark that destabilizes nucleosomes by promoting dissociation of the H2A-H2B dimers from nucleosomes (PubMed: <a href="http://www.uniprot.org/citations/31542297" target=" blank">31542297</a>).

# **Cellular Location**

Nucleus. Chromosome Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Note=Mainly localizes to the nucleus (PubMed:27796307). Localizes to sites of DNA damage (PubMed:25593309) Also localizes to centrosomes in late G1 and around the G1/S transition, coinciding with the onset of centriole formation (PubMed:27796307).

## **Tissue Location**

Expressed in all tissues tested.

### KAT2A Rabbit mAb - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides





• Dot Blot

- <u>Immunohistochemistry</u>
- <u>Immunofluorescence</u>
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

# KAT2A Rabbit mAb - Images

