

**NONO Rabbit mAb**  
**Catalog # AP75813****Specification****NONO Rabbit mAb - Product Information**

Application	WB, IHC-P, IHC-F, ICC
Primary Accession	<a href="#">Q15233</a>
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Monoclonal Antibody
Calculated MW	54232

**NONO Rabbit mAb - Additional Information****Gene ID** 4841**Other Names**  
NONO**Dilution**  
WB~~1/500-1/1000  
IHC-P~~N/A  
IHC-F~~N/A  
ICC~~N/A**Format**  
Liquid**NONO Rabbit mAb - Protein Information****Name** NONO {ECO:0000303|PubMed:9393982, ECO:0000312|HGNC:HGNC:7871}**Function**

DNA- and RNA binding protein, involved in several nuclear processes (PubMed:[11525732](http://www.uniprot.org/citations/11525732), PubMed:[12403470](http://www.uniprot.org/citations/12403470), PubMed:[26571461](http://www.uniprot.org/citations/26571461)). Binds the conventional octamer sequence in double-stranded DNA (PubMed:[11525732](http://www.uniprot.org/citations/11525732), PubMed:[12403470](http://www.uniprot.org/citations/12403470), PubMed:[26571461](http://www.uniprot.org/citations/26571461)). Also binds single- stranded DNA and RNA at a site independent of the duplex site (PubMed:[11525732](http://www.uniprot.org/citations/11525732), PubMed:[12403470](http://www.uniprot.org/citations/12403470), PubMed:[26571461](http://www.uniprot.org/citations/26571461)). Involved in pre- mRNA splicing, probably as a heterodimer with SFPQ (PubMed:[11525732](http://www.uniprot.org/citations/11525732), PubMed:[12403470](http://www.uniprot.org/citations/12403470), PubMed:[26571461](http://www.uniprot.org/citations/26571461)).

[26571461](http://www.uniprot.org/citations/26571461)). Interacts with U5 snRNA, probably by binding to a purine-rich sequence located on the 3' side of U5 snRNA stem 1b (PubMed:[12403470](http://www.uniprot.org/citations/12403470)). Together with PSPC1, required for the formation of nuclear paraspeckles (PubMed:[22416126](http://www.uniprot.org/citations/22416126)). The SFPQ-NONO heteromer associated with MATR3 may play a role in nuclear retention of defective RNAs (PubMed:[11525732](http://www.uniprot.org/citations/11525732)). The SFPQ-NONO heteromer may be involved in DNA unwinding by modulating the function of topoisomerase I/TOP1 (PubMed:[10858305](http://www.uniprot.org/citations/10858305)). The SFPQ-NONO heteromer may be involved in DNA non-homologous end joining (NHEJ) required for double-strand break repair and V(D)J recombination and may stabilize paired DNA ends (PubMed:[15590677](http://www.uniprot.org/citations/15590677)). In vitro, the complex strongly stimulates DNA end joining, binds directly to the DNA substrates and cooperates with the Ku70/G22P1-Ku80/XRCC5 (Ku) dimer to establish a functional preligation complex (PubMed:[15590677](http://www.uniprot.org/citations/15590677)). NONO is involved in transcriptional regulation. The SFPQ-NONO-NR5A1 complex binds to the CYP17 promoter and regulates basal and cAMP-dependent transcriptional activity (PubMed:[11897684](http://www.uniprot.org/citations/11897684)). NONO binds to an enhancer element in long terminal repeats of endogenous intracisternal A particles (IAPs) and activates transcription (By similarity). Regulates the circadian clock by repressing the transcriptional activator activity of the CLOCK-BMAL1 heterodimer (By similarity). Important for the functional organization of GABAergic synapses (By similarity). Plays a specific and important role in the regulation of synaptic RNAs and GPHN/gephyrin scaffold structure, through the regulation of GABRA2 transcript (By similarity). Plays a key role during neuronal differentiation by recruiting TET1 to genomic loci and thereby regulating 5-hydroxymethylcytosine levels (By similarity). Plays a role in the regulation of DNA virus-mediated innate immune response by assembling into the HDP-RNP complex, a complex that serves as a platform for IRF3 phosphorylation and subsequent innate immune response activation through the cGAS-STING pathway (PubMed:[28712728](http://www.uniprot.org/citations/28712728), PubMed:[30270045](http://www.uniprot.org/citations/30270045)). Promotes activation of the cGAS-STING pathway in response to HIV-2 infection: acts by interacting with HIV-2 Capsid protein p24, thereby promoting detection of viral DNA by CGAS, leading to CGAS-mediated immune activation (PubMed:[30270045](http://www.uniprot.org/citations/30270045)). In contrast, the weak interaction with HIV-1 Capsid protein p24 does not allow activation of the cGAS-STING pathway (PubMed:[30270045](http://www.uniprot.org/citations/30270045)).

### Cellular Location

Nucleus. Nucleus, nucleolus. Nucleus speckle. Chromosome {ECO:0000250|UniProtKB:Q99K48}. Note=Detected in punctate subnuclear structures often located adjacent to splicing speckles, called paraspeckles.

### Tissue Location

Heart, brain, placenta, lung, liver, skeletal muscle, kidney and pancreas. Also found in a number of breast tumor cell lines.

## NONO Rabbit mAb - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)

- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### **NONO Rabbit mAb - Images**

