

Rab5 Rabbit mAb
Catalog # AP75987**Specification**

Rab5 Rabbit mAb - Product Information

Application	WB, IHC-P
Primary Accession	P20339
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Monoclonal Antibody
Calculated MW	23659

Rab5 Rabbit mAb - Additional Information**Gene ID** 5868**Other Names**
RAB5A**Dilution**
WB~~1/500-1/1000
IHC-P~~N/A**Format**
50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and 0.05% BSA.**Storage**
Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.**Rab5 Rabbit mAb - Protein Information****Name** RAB5A ([HGNC:9783](#))**Synonyms** RAB5**Function**

The small GTPases Rab are key regulators of intracellular membrane trafficking, from the formation of transport vesicles to their fusion with membranes. Rabs cycle between an inactive GDP-bound form and an active GTP-bound form that is able to recruit to membranes different sets of downstream effectors directly responsible for vesicle formation, movement, tethering and fusion. RAB5A is required for the fusion of plasma membranes and early endosomes and involved in early endocytic trafficking (PubMed:10818110, PubMed:14617813, PubMed:15378032, PubMed:16086013, PubMed:16410077, PubMed:17562788). Required for EEA1 recruitment to early endosomes (PubMed:17562788).

[16086013](http://www.uniprot.org/citations/16086013), PubMed:<[17562788](http://www.uniprot.org/citations/17562788)>). Recruits FERRY complex to early endosomes, where FERRY links early endosomes with a subgroup of mRNAs to enable mRNA intracellular distribution (PubMed:<[37267906](http://www.uniprot.org/citations/37267906)>). Recruits RABEP1/Rabaptin- 5 effector to early endosomes, thereby promoting endocytic membrane fusion (By similarity). Required for EGF and transferrin endocytosis and trafficking through early endosomes (PubMed:<[16086013](http://www.uniprot.org/citations/16086013)>, PubMed:<[17562788](http://www.uniprot.org/citations/17562788)>). Contributes to the regulation of filopodia extension (PubMed:<[14978216](http://www.uniprot.org/citations/14978216)>). Required for the exosomal release of SDCBP, CD63, PDCD6IP and syndecan (PubMed:<[22660413](http://www.uniprot.org/citations/22660413)>). Regulates maturation of apoptotic cell-containing phagosomes, probably downstream of DYN2 and PIK3C3 (By similarity).

Cellular Location

Cell membrane; Lipid-anchor; Cytoplasmic side. Early endosome membrane; Lipid-anchor. Melanosome Cytoplasmic vesicle. Cell projection, ruffle {ECO:0000250|UniProtKB:P18066}. Membrane. Cytoplasm, cytosol. Cytoplasmic vesicle, phagosome membrane {ECO:0000250|UniProtKB:Q9CQD1}. Endosome membrane Note=Enriched in stage I melanosomes (PubMed:17081065). Alternates between membrane-bound and cytosolic forms (Probable) {ECO:0000269|PubMed:17081065, ECO:0000305}

Rab5 Rabbit mAb - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Rab5 Rabbit mAb - Images



