

TFG Rabbit mAb
Catalog # AP76205**Specification**

TFG Rabbit mAb - Product Information

Application	WB, IHC-P, IHC-F, ICC
Primary Accession	Q92734
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Monoclonal Antibody
Calculated MW	43448

TFG Rabbit mAb - Additional Information**Gene ID** 10342**Other Names**
TFG**Dilution**
WB~~1/500-1/1000
IHC-P~~N/A
IHC-F~~N/A
ICC~~N/A**Format**
Liquid**TFG Rabbit mAb - Protein Information****Name** TFG**Function**

Plays a role in the normal dynamic function of the endoplasmic reticulum (ER) and its associated microtubules (PubMed:<[a href="http://www.uniprot.org/citations/23479643" target="_blank">23479643](http://www.uniprot.org/citations/23479643)>, PubMed:<[a href="http://www.uniprot.org/citations/27813252" target="_blank">27813252](http://www.uniprot.org/citations/27813252)>). Required for secretory cargo traffic from the endoplasmic reticulum to the Golgi apparatus (PubMed:<[a href="http://www.uniprot.org/citations/21478858" target="_blank">21478858](http://www.uniprot.org/citations/21478858)>).

Cellular Location

Endoplasmic reticulum. Note=Localizes to endoplasmic reticulum exit site (ERES), also known as transitional endoplasmic reticulum (tER) (PubMed:21478858, PubMed:27813252).

Tissue Location

Ubiquitous.

TFG Rabbit mAb - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

TFG Rabbit mAb - Images



