

Phospho-Elongation Factor 2 (Thr56/Thr58) Rabbit mAb
Catalog # AP76334**Specification****Phospho-Elongation Factor 2 (Thr56/Thr58) Rabbit mAb - Product Information**

Application	WB
Primary Accession	P13639
Reactivity	Human
Host	Rabbit
Clonality	Monoclonal Antibody

Phospho-Elongation Factor 2 (Thr56/Thr58) Rabbit mAb - Additional Information**Gene ID** 1938**Other Names**
EEF2**Dilution**
WB~~1/500-1/1000**Format**
Liquid**Phospho-Elongation Factor 2 (Thr56/Thr58) Rabbit mAb - Protein Information****Name** EEF2**Synonyms** EF2**Function**

Catalyzes the GTP-dependent ribosomal translocation step during translation elongation (PubMed: [26593721](http://www.uniprot.org/citations/26593721)). During this step, the ribosome changes from the pre-translocational (PRE) to the post-translocational (POST) state as the newly formed A-site-bound peptidyl- tRNA and P-site-bound deacylated tRNA move to the P and E sites, respectively (PubMed: [26593721](http://www.uniprot.org/citations/26593721)). Catalyzes the coordinated movement of the two tRNA molecules, the mRNA and conformational changes in the ribosome (PubMed: [26593721](http://www.uniprot.org/citations/26593721)).

Cellular Location

Cytoplasm. Nucleus. Note=Phosphorylation by CSK promotes cleavage and SUMOylation-dependent nuclear translocation of the C- terminal cleavage product.

Phospho-Elongation Factor 2 (Thr56/Thr58) Rabbit mAb - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Phospho-Elongation Factor 2 (Thr56/Thr58) Rabbit mAb - Images

