

BUBR1 Rabbit mAb
Catalog # AP76412**Specification**

BUBR1 Rabbit mAb - Product Information

Application	WB, IP, ICC
Primary Accession	O60566
Reactivity	Human
Host	Rabbit
Clonality	Monoclonal Antibody
Calculated MW	119545

BUBR1 Rabbit mAb - Additional Information**Gene ID** 701**Other Names**
BUB1B**Dilution**
WB~~1/500-1/1000
IP~~N/A
ICC~~N/A**Format**
Liquid**BUBR1 Rabbit mAb - Protein Information****Name** BUB1B**Synonyms** BUBR1, MAD3L, SSK1**Function**

Essential component of the mitotic checkpoint. Required for normal mitosis progression. The mitotic checkpoint delays anaphase until all chromosomes are properly attached to the mitotic spindle. One of its checkpoint functions is to inhibit the activity of the anaphase-promoting complex/cyclosome (APC/C) by blocking the binding of CDC20 to APC/C, independently of its kinase activity. The other is to monitor kinetochore activities that depend on the kinetochore motor CENPE. Required for kinetochore localization of CENPE. Negatively regulates PLK1 activity in interphase cells and suppresses centrosome amplification. Also implicated in triggering apoptosis in polyploid cells that exit aberrantly from mitotic arrest. May play a role for tumor suppression.

Cellular Location

Cytoplasm. Nucleus. Chromosome, centromere, kinetochore. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Note=Cytoplasmic in interphase cells. Associates with the kinetochores in early prophase. Kinetochore localization requires BUB1, PLK1 and KNL1

Tissue Location

Highly expressed in thymus followed by spleen. Preferentially expressed in tissues with a high mitotic index

BUBR1 Rabbit mAb - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

BUBR1 Rabbit mAb - Images

