

**Dnmt3a Rabbit mAb**  
**Catalog # AP76472****Specification****Dnmt3a Rabbit mAb - Product Information**

Application	WB
Primary Accession	<a href="#">Q9Y6K1</a>
Reactivity	Human
Host	Rabbit
Clonality	Monoclonal Antibody
Calculated MW	101858

**Dnmt3a Rabbit mAb - Additional Information****Gene ID** 1788**Other Names**

DNMT3A

**Dilution**

WB~~1/500-1/1000

**Format**

50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and 0.05% BSA.

**Storage**

Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

**Dnmt3a Rabbit mAb - Protein Information****Name** DNMT3A**Function**

Required for genome-wide de novo methylation and is essential for the establishment of DNA methylation patterns during development (PubMed:<a href="http://www.uniprot.org/citations/12138111" target="\_blank">12138111</a>, PubMed:<a href="http://www.uniprot.org/citations/16357870" target="\_blank">16357870</a>, PubMed:<a href="http://www.uniprot.org/citations/30478443" target="\_blank">30478443</a>). DNA methylation is coordinated with methylation of histones (PubMed:<a href="http://www.uniprot.org/citations/12138111" target="\_blank">12138111</a>, PubMed:<a href="http://www.uniprot.org/citations/16357870" target="\_blank">16357870</a>, PubMed:<a href="http://www.uniprot.org/citations/30478443" target="\_blank">30478443</a>). It modifies DNA in a non-processive manner and also methylates non-CpG sites (PubMed:<a href="http://www.uniprot.org/citations/12138111" target="\_blank">12138111</a>, PubMed:<a href="http://www.uniprot.org/citations/16357870" target="\_blank">16357870</a>, PubMed:<a href="http://www.uniprot.org/citations/30478443" target="\_blank">30478443</a>). May preferentially methylate DNA linker between 2 nucleosomal cores and is inhibited by histone H1 (By similarity). Plays a role in paternal and maternal imprinting (By similarity). Required for

methylation of most imprinted loci in germ cells (By similarity). Acts as a transcriptional corepressor for ZBTB18 (By similarity). Recruited to trimethylated 'Lys-36' of histone H3 (H3K36me3) sites (By similarity). Can actively repress transcription through the recruitment of HDAC activity (By similarity). Also has weak auto-methylation activity on Cys-710 in absence of DNA (By similarity).

#### Cellular Location

Nucleus. Chromosome Cytoplasm. Note=Accumulates in the major satellite repeats at pericentric heterochromatin {ECO:0000250|UniProtKB:O88508}

#### Tissue Location

Highly expressed in fetal tissues, skeletal muscle, heart, peripheral blood mononuclear cells, kidney, and at lower levels in placenta, brain, liver, colon, spleen, small intestine and lung

### Dnmt3a Rabbit mAb - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### Dnmt3a Rabbit mAb - Images

