

Elongation factor 2 Rabbit mAb
Catalog # AP76477**Specification**

Elongation factor 2 Rabbit mAb - Product Information

Application	WB, IHC-P, IHC-F, IP, ICC
Primary Accession	P13639
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Monoclonal Antibody
Calculated MW	95338

Elongation factor 2 Rabbit mAb - Additional Information**Gene ID** 1938**Other Names**
EEF2**Dilution**
WB~~1/500-1/1000
IHC-P~~N/A
IHC-F~~N/A
IP~~1/20
ICC~~N/A**Format**
50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and 0.05% BSA.**Elongation factor 2 Rabbit mAb - Protein Information****Name** EEF2**Synonyms** EF2**Function**

Catalyzes the GTP-dependent ribosomal translocation step during translation elongation (PubMed: [26593721](http://www.uniprot.org/citations/26593721)). During this step, the ribosome changes from the pre-translocational (PRE) to the post-translocational (POST) state as the newly formed A-site-bound peptidyl- tRNA and P-site-bound deacylated tRNA move to the P and E sites, respectively (PubMed: [26593721](http://www.uniprot.org/citations/26593721)). Catalyzes the coordinated movement of the two tRNA molecules, the mRNA and conformational changes in the ribosome (PubMed: [26593721](http://www.uniprot.org/citations/26593721)).

Cellular Location

Cytoplasm. Nucleus. Note=Phosphorylation by CSK promotes cleavage and

SUMOylation-dependent nuclear translocation of the C- terminal cleavage product.

Elongation factor 2 Rabbit mAb - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Elongation factor 2 Rabbit mAb - Images



