

**CREB Regulated Transcription Coactivator 2 Rabbit mAb**  
**Catalog # AP76743****Specification**

---

**CREB Regulated Transcription Coactivator 2 Rabbit mAb - Product Information**

Application	WB, IHC-P, IP
Primary Accession	<a href="#">Q53ET0</a>
Reactivity	Human
Host	Rabbit
Clonality	Monoclonal Antibody
Calculated MW	73302

**CREB Regulated Transcription Coactivator 2 Rabbit mAb - Additional Information****Gene ID** 200186**Other Names**  
CRTC2**Dilution**  
WB~~1/500-1/1000  
IHC-P~~N/A  
IP~~N/A**Format**  
Liquid**CREB Regulated Transcription Coactivator 2 Rabbit mAb - Protein Information****Name** CRTC2**Synonyms** TORC2**Function**

Transcriptional coactivator for CREB1 which activates transcription through both consensus and variant cAMP response element (CRE) sites. Acts as a coactivator, in the SIK/TORC signaling pathway, being active when dephosphorylated and acts independently of CREB1 'Ser-133' phosphorylation. Enhances the interaction of CREB1 with TAF4. Regulates gluconeogenesis as a component of the LKB1/AMPK/TORC2 signaling pathway. Regulates the expression of specific genes such as the steroidogenic gene, StAR. Potent coactivator of PPARGC1A and inducer of mitochondrial biogenesis in muscle cells. Also coactivator for TAX activation of the human T-cell leukemia virus type 1 (HTLV-1) long terminal repeats (LTR).

**Cellular Location**

Cytoplasm. Nucleus. Note=Translocated from the nucleus to the cytoplasm on interaction of the phosphorylated form with 14-3-3 protein (PubMed:15454081). In response to cAMP levels and glucagon, relocated to the nucleus (PubMed:15454081)

### Tissue Location

Most abundantly expressed in the thymus. Present in both B and T-lymphocytes. Highly expressed in HEK293T cells and in insulinomas. High levels also in spleen, ovary, muscle and lung, with highest levels in muscle. Lower levels found in brain, colon, heart, kidney, prostate, small intestine and stomach. Weak expression in liver and pancreas.

### CREB Regulated Transcription Coactivator 2 Rabbit mAb - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### CREB Regulated Transcription Coactivator 2 Rabbit mAb - Images



