

DPP3 Rabbit mAb
Catalog # AP78787**Specification**

DPP3 Rabbit mAb - Product Information

Application	WB, FC
Primary Accession	Q9NY33
Reactivity	Human
Host	Rabbit
Clonality	Monoclonal Antibody
Calculated MW	82589

DPP3 Rabbit mAb - Additional Information**Gene ID** 10072**Other Names**
DPP3**Dilution**
WB~~1/500-1/1000
FC~~1:10~50**Format**
Liquid**DPP3 Rabbit mAb - Protein Information****Name** DPP3**Function**

Cleaves and degrades bioactive peptides, including angiotensin, Leu-enkephalin and Met-enkephalin (PubMed:<[a href="http://www.uniprot.org/citations/1515063" target="_blank">1515063](http://www.uniprot.org/citations/1515063), PubMed:<[a href="http://www.uniprot.org/citations/3233187" target="_blank">3233187](http://www.uniprot.org/citations/3233187)). Also cleaves Arg-Arg-beta-naphthylamide (in vitro) (PubMed:<[a href="http://www.uniprot.org/citations/11209758" target="_blank">11209758](http://www.uniprot.org/citations/11209758), PubMed:<[a href="http://www.uniprot.org/citations/3233187" target="_blank">3233187](http://www.uniprot.org/citations/3233187), PubMed:<[a href="http://www.uniprot.org/citations/9425109" target="_blank">9425109](http://www.uniprot.org/citations/9425109)).

Cellular Location
Cytoplasm, cytosol**Tissue Location**

Detected in placenta (at protein level) (PubMed:3233187). Detected in erythrocytes (at protein level) (PubMed:1515063).

DPP3 Rabbit mAb - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

DPP3 Rabbit mAb - Images

