

cGKI (cGKI beta) Antibody (C-term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP8000a**Specification**

cGKI (cGKI beta) Antibody (C-term) - Product Information

Application	IHC-P, WB,E
Primary Accession	Q13976
Other Accession	P00516 , P14619
Reactivity	Human, Mouse
Predicted	Bovine
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Antigen Region	629-660

cGKI (cGKI beta) Antibody (C-term) - Additional Information**Gene ID** 5592**Other Names**

cGMP-dependent protein kinase 1, cGK 1, cGK1, cGMP-dependent protein kinase I, cGKI, PRKG1, PRKG1B, PRKGR1A, PRKGR1B

Target/Specificity

This cGKI (cGKI beta) antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 629-660 amino acids from the C-terminal region of human cGKI beta.

Dilution

IHC-P~~1:100

WB~~1:1000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

cGKI (cGKI beta) Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

cGKI (cGKI beta) Antibody (C-term) - Protein Information**Name** PRKG1

Synonyms PRKG1B, PRKGR1A, PRKGR1B

Function Serine/threonine protein kinase that acts as a key mediator of the nitric oxide (NO)/cGMP signaling pathway. GMP binding activates PRKG1, which phosphorylates serines and threonines on many cellular proteins. Numerous protein targets for PRKG1 phosphorylation are implicated in modulating cellular calcium, but the contribution of each of these targets may vary substantially among cell types. Proteins that are phosphorylated by PRKG1 regulate platelet activation and adhesion, smooth muscle contraction, cardiac function, gene expression, feedback of the NO-signaling pathway, and other processes involved in several aspects of the CNS like axon guidance, hippocampal and cerebellar learning, circadian rhythm and nociception. Smooth muscle relaxation is mediated through lowering of intracellular free calcium, by desensitization of contractile proteins to calcium, and by decrease in the contractile state of smooth muscle or in platelet activation. Regulates intracellular calcium levels via several pathways: phosphorylates IRAG1 and inhibits IP3-induced Ca(2+) release from intracellular stores, phosphorylation of KCNMA1 (BKCa) channels decreases intracellular Ca(2+) levels, which leads to increased opening of this channel. PRKG1 phosphorylates the canonical transient receptor potential channel (TRPC) family which inactivates the associated inward calcium current. Another mode of action of NO/cGMP/PKG1 signaling involves PKGI-mediated inactivation of the Ras homolog gene family member A (RhoA). Phosphorylation of RHOA by PRKG1 blocks the action of this protein in myriad processes: regulation of RHOA translocation; decreasing contraction; controlling vesicle trafficking, reduction of myosin light chain phosphorylation resulting in vasorelaxation. Activation of PRKG1 by NO signaling also alters gene expression in a number of tissues. In smooth muscle cells, increased cGMP and PRKG1 activity influence expression of smooth muscle-specific contractile proteins, levels of proteins in the NO/cGMP signaling pathway, down- regulation of the matrix proteins osteopontin and thrombospondin-1 to limit smooth muscle cell migration and phenotype. Regulates vasodilator-stimulated phosphoprotein (VASP) functions in platelets and smooth muscle.

Cellular Location

Cytoplasm. Note=Colocalized with TRPC7 in the plasma membrane.


Tissue Location


Primarily expressed in lung and placenta.

cGKI (cGKI beta) Antibody (C-term) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


cGKI (cGKI beta) Antibody (C-term) - Images




All lanes : Anti-cGKI (cGKI beta) Antibody (C-term) at 1:1000 dilution Lane 1: 293 whole cell lysate Lane 2: Hela whole cell lysate Lane 3: U-87 MG whole cell lysate Lane 4: Mouse small intestine tissue lysate Lane 5: NIH/3T3 whole cell lysate Lane 6: C6 whole cell lysate Lysates/proteins at 20 μ g per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 76 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



Anti-cGKI (cGKI beta) Antibody (C-term) at 1:1000 dilution + SH-SY5Y whole cell lysate Lysates/proteins at 20 μ g per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 76 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



Immunohistochemical analysis of AP8000A on paraffin-embedded Human placenta tissue. Tissue was fixed with formaldehyde at room temperature. Heat induced epitope retrieval was performed by EDTA buffer (pH9. 0). Samples were incubated with primary antibody(1:100) for 1 hour at room temperature. Undiluted CRF Anti-Polyvalent HRP Polymer antibody was used as the secondary antibody.



Immunohistochemical analysis of AP8000A on paraffin-embedded Human hepato carcinoma tissue. Tissue was fixed with formaldehyde at room temperature. Heat induced epitope retrieval was performed by EDTA buffer (pH9. 0). Samples were incubated with primary antibody(1:100) for 1 hour at room temperature. Undiluted CRF Anti-Polyvalent HRP Polymer antibody was used as the secondary antibody.

cGKI (cGKI beta) Antibody (C-term) - Background

Protein kinases are enzymes that transfer a phosphate group from a phosphate donor, generally the γ phosphate of ATP, onto an acceptor amino acid in a substrate protein. By this basic mechanism, protein kinases mediate most of the signal transduction in eukaryotic cells, regulating cellular metabolism, transcription, cell cycle progression, cytoskeletal rearrangement and cell movement, apoptosis, and differentiation. With more than 500 gene products, the protein kinase family is one of the largest families of proteins in eukaryotes. The family has been classified in 8 major groups based on sequence comparison of their tyrosine (PTK) or serine/threonine (STK) kinase catalytic domains.

cGKI (cGKI beta) Antibody (C-term) - References

Orstavik, S., et al., Genomics 42(2):311-318 (1997).

Tamura, N., et al., Hypertension 27 (3 Pt 2), 552-557 (1996).

cGKI (cGKI beta) Antibody (C-term) - Citations

- [A molecular mechanism for therapeutic effects of cGMP-elevating agents in pulmonary arterial hypertension.](#)