

**PCK1 Antibody (N-term)**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP8093a**

**Specification**

---

**PCK1 Antibody (N-term) - Product Information**

|                   |                        |
|-------------------|------------------------|
| Application       | WB, IHC-P,E            |
| Primary Accession | <a href="#">P35558</a> |
| Reactivity        | Human, Mouse           |
| Host              | Rabbit                 |
| Clonality         | Polyclonal             |
| Isotype           | Rabbit IgG             |
| Calculated MW     | 69195                  |
| Antigen Region    | 40-70                  |

**PCK1 Antibody (N-term) - Additional Information**

**Gene ID** 5105

**Other Names**

Phosphoenolpyruvate carboxykinase, cytosolic [GTP], PEPCK-C, PCK1, PEPCK1

**Target/Specificity**

This PCK1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 40-70 amino acids from the N-terminal region of human PCK1.

**Dilution**

WB~~1:1000

IHC-P~~1:50~100

E~~Use at an assay dependent concentration.

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

PCK1 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

**PCK1 Antibody (N-term) - Protein Information**

**Name** PCK1 {ECO:0000303|PubMed:8490617, ECO:0000312|HGNC:HGNC:8724}

**Function** Cytosolic phosphoenolpyruvate carboxykinase that catalyzes the reversible

decarboxylation and phosphorylation of oxaloacetate (OAA) and acts as the rate-limiting enzyme in gluconeogenesis (PubMed:[24863970](#), PubMed:[26971250](#), PubMed:[28216384](#), PubMed:[30193097](#)). Regulates cataplerosis and anaplerosis, the processes that control the levels of metabolic intermediates in the citric acid cycle (PubMed:[24863970](#), PubMed:[26971250](#), PubMed:[28216384](#), PubMed:[30193097](#)). At low glucose levels, it catalyzes the cataplerotic conversion of oxaloacetate to phosphoenolpyruvate (PEP), the rate-limiting step in the metabolic pathway that produces glucose from lactate and other precursors derived from the citric acid cycle (PubMed:[30193097](#)). At high glucose levels, it catalyzes the anaplerotic conversion of phosphoenolpyruvate to oxaloacetate (PubMed:[30193097](#)). Acts as a regulator of formation and maintenance of memory CD8(+) T-cells: up- regulated in these cells, where it generates phosphoenolpyruvate, via gluconeogenesis (By similarity). The resultant phosphoenolpyruvate flows to glycogen and pentose phosphate pathway, which is essential for memory CD8(+) T-cells homeostasis (By similarity). In addition to the phosphoenolpyruvate carboxykinase activity, also acts as a protein kinase when phosphorylated at Ser-90: phosphorylation at Ser-90 by AKT1 reduces the binding affinity to oxaloacetate and promotes an atypical serine protein kinase activity using GTP as donor (PubMed:[32322062](#)). The protein kinase activity regulates lipogenesis: upon phosphorylation at Ser-90, translocates to the endoplasmic reticulum and catalyzes phosphorylation of INSIG proteins (INSIG1 and INSIG2), thereby disrupting the interaction between INSIG proteins and SCAP and promoting nuclear translocation of SREBP proteins (SREBF1/SREBP1 or SREBF2/SREBP2) and subsequent transcription of downstream lipogenesis- related genes (PubMed:[32322062](#)).

#### Cellular Location

Cytoplasm, cytosol. Endoplasmic reticulum Note=Phosphorylation at Ser-90 promotes translocation to the endoplasmic reticulum.

#### Tissue Location

Major sites of expression are liver, kidney and adipocytes.


#### PCK1 Antibody (N-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### PCK1 Antibody (N-term) - Images






The anti-PCK1 Pab (Cat. #AP8093a) is used in Western blot to detect PCK1 in mouse liver tissue lysate.



PCK1 Antibody (N-term) (Cat. #AP8093a) western blot analysis in MCF-7 cell line lysates (35ug/lane). This demonstrates the PCK1 antibody detected the PCK1 protein (arrow).



Formalin-fixed and paraffin-embedded human cancer tissue reacted with the primary antibody, which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated. BC = breast carcinoma; HC = hepatocarcinoma.

#### **PCK1 Antibody (N-term) - Background**

This gene is a main control point for the regulation of gluconeogenesis. The cytosolic enzyme encoded by this gene, along with GTP, catalyzes the formation of phosphoenolpyruvate from oxaloacetate, with the release of carbon dioxide and GDP. The expression of this gene can be regulated by insulin, glucocorticoids, glucagon, cAMP, and diet. A mitochondrial isozyme of the encoded protein also has been characterized.

#### **PCK1 Antibody (N-term) - References**

- Berger, K. et al. PLoS ONE. 4(3): e4671 (2009).  
Dunten, P., et al., J. Mol. Biol. 316(2):257-264 (2002).  
Strausberg, R.L., et al., Proc. Natl. Acad. Sci. U.S.A. 99(26):16899-16903 (2002).  
Deloukas, P., et al., Nature 414(6866):865-871 (2001).  
O'Brien, R.M., et al., Biochim. Biophys. Acta 1264(3):284-288 (1995).  
Ting, C.N., et al., Genomics 16(3):698-706 (1993).